

# *Selfextensional logics with a distributive nearlattice term*

**Luciano J. González**

**Archive for Mathematical Logic**


ISSN 0933-5846

Volume 58

Combined 1-2

Arch. Math. Logic (2019) 58:219-243

DOI 10.1007/s00153-018-0628-1



**Your article is protected by copyright and all rights are held exclusively by Springer-Verlag GmbH Germany, part of Springer Nature. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**



# Selfextensional logics with a distributive nearlattice term

Luciano J. González<sup>1</sup> 

Received: 31 January 2018 / Accepted: 2 May 2018 / Published online: 11 May 2018  
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

**Abstract** We define when a ternary term  $m$  of an algebraic language  $\mathcal{L}$  is called a *distributive nearlattice term* (DN-term) of a sentential logic  $\mathcal{S}$ . Distributive nearlattices are ternary algebras generalising Tarski algebras and distributive lattices. We characterise the selfextensional logics with a DN-term through the interpretation of the DN-term in the algebras of the algebraic counterpart of the logics. We prove that the canonical class of algebras (under the point of view of Abstract Algebraic Logic) associated with a selfextensional logic with a DN-term is a variety, and we obtain that the logic is in fact fully selfextensional.

**Keywords** Selfextensional logic · Distributive nearlattice · Logics based on partial orders · Abstract algebraic logic

**Mathematics Subject Classification** 03G27 · 03B22 · 03G25 · 06A12

## 1 Introduction

This paper is motivated by the results and ideas given in [33, 34]. In [33], selfextensional finitary logics with a binary term  $\rightarrow$  satisfying the deduction-detachment theorem are studied. There, these logics are characterised as logics  $\mathcal{S}$  for which there is a class

---

This work was partially supported by Universidad Nacional de La Pampa (Facultad de Ciencias Exactas y Naturales) under the Grant P.I. 64 M, Res. 432/14 CD; and also by Consejo Nacional de Investigaciones Científicas y Técnicas (Argentina) under the Grant PIP 112-20150-100412CO.

---

✉ Luciano J. González  
lucianogonzalez@exactas.unlpam.edu.ar

<sup>1</sup> Facultad de Ciencias Exactas y Naturales, Universidad Nacional de La Pampa, Santa Rosa, Argentina

of algebras  $K$  such that the equations defining Hilbert algebras (also called positive implication algebras) hold for  $\rightarrow$  and the following condition is satisfied:

$$\varphi_0, \dots, \varphi_{n-1} \vdash_{\mathcal{S}} \varphi \iff (\forall A \in K)(\forall h \in \text{Hom}(Fm, A)) \\ h(\varphi_0 \rightarrow (\dots \rightarrow (\varphi_{n-1} \rightarrow \varphi) \dots)) = 1.$$

Similar results are obtained in [34]. There, selfextensional finitary logics with a conjunction term  $\wedge$  are characterised as the logics  $\mathcal{S}$  for which there is a class of algebras  $K$  such that the semilattice equations are satisfied for  $\wedge$  and the following condition holds:

$$\varphi_0, \dots, \varphi_{n-1} \vdash_{\mathcal{S}} \varphi \iff (\forall A \in K)(\forall h \in \text{Hom}(Fm, A)) \\ h(\varphi_0) \wedge \dots \wedge h(\varphi_{n-1}) \leq h(\varphi).$$

As we can notice from the definitions above, the two kinds of sentential logic are characterised through the behaviour of the interpretation of the implication  $\rightarrow$  and the conjunction  $\wedge$  in the corresponding classes of algebras  $K$ .

The notion of *distributive nearlattice* can be presented in two different and equivalent ways. They can be defined as join-semilattices with some extra properties and can be defined as algebras with only one ternary operation satisfying some identities. The two different ways to consider distributive nearlattices are useful for various purposes. After the formal definition of distributive nearlattice (Definition 2.5), we will see (Remark 2.10) that the variety of distributive nearlattices is a natural generalisation of both the variety of Tarski algebras (also called implication algebras) and the variety of distributive lattices.

The primary aim of this paper is to propose a definition of when a ternary term  $m$  of an algebraic language  $\mathcal{L}$  can be considered a *distributive nearlattice term* (*DN-term* for short) for a sentential logic  $\mathcal{S}$ . We present some syntactical properties (Sect. 3) on a sentential logic  $\mathcal{S}$  concerning a ternary term  $m$  such that, when  $m$  is interpreted in every algebra  $A$  of the algebraic counterpart of the logic  $\mathcal{S}$ , the  $\{m\}$ -reduct  $\langle A, m^A \rangle$  will be a distributive nearlattice.

We show that selfextensional logics with a distributive nearlattice term  $m$  can be characterised as logics  $\mathcal{S}$  for which there exists a class of algebras  $K$  such that the  $\{m\}$ -reducts of the algebras of  $K$  are distributive nearlattices and the consequence relation of  $\mathcal{S}$  can be defined using the partial order induced by the term  $m$  on the algebras of  $K$  (Sect. 3).

In Sect. 5, given a selfextensional logic  $\mathcal{S}$  with a DN-term (and with theorems), we consider two sentential logics associated with the canonical class of algebras of  $\mathcal{S}$ ; namely, *the logic preserving degrees of truth* and *the logic preserving truth*. We show some properties of these logics, and we present some sufficient conditions for these logics to coincide with the original logic  $\mathcal{S}$ .

## 2 Preliminaries

In this section, we introduce some basic concepts and results of Abstract Algebraic Logic (AAL) needed for what follows in the paper and we present the algebraic theory

of nearlattices. Our main references for AAL are [17,23,26,27] and for the theory of nearlattice are [10,13,32].

### 2.1 Abstract algebraic logic

Let  $\mathcal{L}$  be an algebraic language (or algebraic similarity type). We denote by  $Fm(\mathcal{L})$  the absolutely free algebra of type  $\mathcal{L}$  with a denumerable set  $Var$  of propositional variables as the set of generators. The algebra  $Fm(\mathcal{L})$  is called the *algebra of formulas* of type  $\mathcal{L}$  and its elements are called *formulas*. When there is no danger of confusion, we write  $Fm$  instead of  $Fm(\mathcal{L})$ .

A *sentential logic* (also called *deductive system* in AAL)<sup>1</sup> of type  $\mathcal{L}$  is a pair  $\mathcal{S} = \langle Fm, \vdash_{\mathcal{S}} \rangle$  where  $Fm$  is the algebra of formulas of type  $\mathcal{L}$  and  $\vdash_{\mathcal{S}} \subseteq \mathcal{P}(Fm) \times Fm$  is a relation satisfying the following properties: for all  $\Gamma, \Delta \subseteq Fm$  and  $\varphi \in Fm$  (as usual we write  $\Gamma \vdash_{\mathcal{S}} \varphi$  for  $(\Gamma, \varphi) \in \vdash_{\mathcal{S}}$ ),

- (S1) if  $\varphi \in \Gamma$ , then  $\Gamma \vdash_{\mathcal{S}} \varphi$ ;
- (S2) if  $\Gamma \vdash_{\mathcal{S}} \varphi$  and  $\Gamma \subseteq \Delta$ , then  $\Delta \vdash_{\mathcal{S}} \varphi$ ;
- (S3) if  $\Gamma \vdash_{\mathcal{S}} \varphi$  and for every  $\gamma \in \Gamma$ ,  $\Delta \vdash_{\mathcal{S}} \gamma$ , then  $\Delta \vdash_{\mathcal{S}} \varphi$ ;
- (S4) if  $\Gamma \vdash_{\mathcal{S}} \varphi$ , then there is a finite  $\Gamma_0 \subseteq \Gamma$  such that  $\Gamma_0 \vdash_{\mathcal{S}} \varphi$ ;
- (S5) if  $\Gamma \vdash_{\mathcal{S}} \varphi$ , then  $\sigma[\Gamma] \vdash_{\mathcal{S}} \sigma(\varphi)$  for all substitution  $\sigma \in \text{Hom}(Fm, Fm)$ .

The relation  $\vdash_{\mathcal{S}}$  is called the *consequence relation* of  $\mathcal{S}$ . A set  $\Gamma \subseteq Fm$  is called a *theory* of  $\mathcal{S}$  ( *$\mathcal{S}$ -theory*, for short) if is closed under the consequence relation of  $\mathcal{S}$ , that is, for every formula  $\varphi \in Fm$ , if  $\Gamma \vdash_{\mathcal{S}} \varphi$ , then  $\varphi \in \Gamma$ . Let us denote by  $\text{Th}(\mathcal{S})$  the collection of all  $\mathcal{S}$ -theories. It is easy to see that  $\text{Th}(\mathcal{S})$  is an algebraic closure system on  $Fm$  and the closure operator associated with  $\text{Th}(\mathcal{S})$ , which is denoted by  $C_{\mathcal{S}}$ , is defined as:

$$\varphi \in C_{\mathcal{S}}(\Gamma) \iff \Gamma \vdash_{\mathcal{S}} \varphi$$

for all  $\Gamma \cup \{\varphi\} \subseteq Fm$ . Moreover, it is clear that  $C_{\mathcal{S}}$  is finitary.

Let  $\mathcal{S}$  be a sentential logic. The *Frege relation* of  $\mathcal{S}$ , in symbols  $\Lambda(\mathcal{S})$ , is the interderivability relation, that is,  $(\varphi, \psi) \in \Lambda(\mathcal{S})$  if and only if  $\varphi \vdash_{\mathcal{S}} \psi$  and  $\psi \vdash_{\mathcal{S}} \varphi$ . The Frege relation of a sentential logic is an equivalence relation but it is not necessarily a congruence on  $Fm$ . A sentential logic  $\mathcal{S}$  is said to be *selfextensional* (or  $\mathcal{S}$  has the *congruence property*) if the Frege relation  $\Lambda(\mathcal{S})$  is a congruence on  $Fm$ .

Let  $A$  be an algebra of the same similarity type as  $\mathcal{S}$ . A subset  $F \subseteq A$  is said to be an  *$\mathcal{S}$ -filter* of  $A$  if and only if for any  $\Gamma \cup \{\varphi\} \subseteq Fm$  and any interpretation  $h \in \text{Hom}(Fm, A)$ ,

$$\text{if } \Gamma \vdash_{\mathcal{S}} \varphi \text{ and } h[\Gamma] \subseteq F, \text{ then } h(\varphi) \in F.$$

<sup>1</sup> Conditions (S1), (S2), (S3) and (S5) define the notion of sentential logic that is usually considered in abstract algebraic logic. The concept of logic considered in this paper is that of finitary sentential logic, that is, sentential logics satisfying (S4).

The set of all  $\mathcal{S}$ -filters on a given algebra  $A$  is denoted by  $\text{Fi}_{\mathcal{S}}(A)$ ; this set is an algebraic closure system. The associated closure operator will be denoted by  $\text{Fi}_{\mathcal{S}}^A$ .

Let  $\mathcal{L}$  be a fixed but arbitrary algebraic language. A *generalized matrix*, *g-matrix* for short, of similarity type  $\mathcal{L}$  is a pair  $\langle A, \mathcal{C} \rangle$  where  $A$  is an algebra of type  $\mathcal{L}$  and  $\mathcal{C}$  is an algebraic closure system on  $A$ . We denote by  $C$  the closure operator associated with  $\mathcal{C}$  and we will often identify the g-matrix  $\langle A, \mathcal{C} \rangle$  with the pair  $\langle A, C \rangle$ . Notice that the closure operator  $C$  is finitary, i.e., for all  $X \cup \{a\} \subseteq A$ ,  $a \in C(X)$  implies that there is a finite  $X_0 \subseteq X$  such that  $a \in C(X_0)$ .

The reader should keep in mind that all logics and g-matrices considered in this paper are finitary and thus some general results of AAL are restricted to this assumptions.

One of the most interesting aspects of g-matrices is that they can be used in a completely natural way both as models of sentential logics and as models of Gentzen systems. This double function of g-matrices allows relating the algebraic theory of sentential logics to the Gentzen systems. We address the interested reader on these topics to [26,27].

An important example of g-matrix is given by a sentential logic  $\mathcal{S}$ . If  $\mathcal{S}$  is a sentential logic, then  $\langle \text{Fm}, \text{Th}(\mathcal{S}) \rangle$  is a g-matrix.

**Definition 2.1** A g-matrix  $\langle A, C \rangle$  is said to be a *g-model* of a sentential logic  $\mathcal{S}$  when for all  $\Gamma \cup \{\varphi\} \subseteq \text{Fm}$ , if  $\Gamma \vdash_{\mathcal{S}} \varphi$ , then  $h(\varphi) \in C(h[\Gamma])$  for all  $h \in \text{Hom}(\text{Fm}, A)$ . Let us denote the class of all g-models of a sentential logic  $\mathcal{S}$  by  $\mathbf{GMod}(\mathcal{S})$ .

The logical concept of Frege relation is transferred to the setting of g-matrices. The *Frege relation* of a g-matrix  $\langle A, C \rangle$  is defined by:

$$(a, b) \in \Lambda_A(C) \iff C(a) = C(b)$$

for every  $a, b \in A$ . The *Tarski congruence* of a g-matrix  $\langle A, C \rangle$  is the largest congruence below the Frege relation of the g-matrix. We denote the Tarski congruence of  $\langle A, C \rangle$  by  $\tilde{\Omega}_A(C)$ . A g-matrix is said to be *reduced* when its Tarski congruence is the identity relation. Let us denote by  $\mathbf{GMod}^*(\mathcal{S})$  the class of all reduced g-models of a sentential logic  $\mathcal{S}$ .

We can now introduce the class of algebras that is considered in AAL as the natural algebraic counterpart of a sentential logic, see [18,27].

**Definition 2.2** The *canonical class of algebras* associated with a sentential logic  $\mathcal{S}$  (it is also called the *algebraic counterpart* of  $\mathcal{S}$ ) is the class of the algebraic reducts of the reduced g-models of  $\mathcal{S}$ ; it is denoted by  $\text{Alg}(\mathcal{S})$ . That is,

$$\begin{aligned} \text{Alg}(\mathcal{S}) &:= \text{Alg}(\mathbf{GMod}^*(\mathcal{S})) \\ &= \{A : \langle A, C \rangle \in \mathbf{GMod}^*(\mathcal{S}) \text{ for some finitary closure operator } C\}. \end{aligned}$$

Moreover, another important class of algebras associated with a sentential logic  $\mathcal{S}$  is  $\mathbf{K}_{\mathcal{S}} := \mathbb{V}(\text{Fm}/\tilde{\Omega}(\mathcal{S}))$ , the variety generated by the algebra  $\text{Fm}/\tilde{\Omega}(\mathcal{S})$ . This variety is called *the intrinsic variety of  $\mathcal{S}$* .

**Lemma 2.3** [26, Proposition 2.26] *Let  $\mathcal{S}$  be a sentential logic. Then, the intrinsic variety of  $\mathcal{S}$  is the variety generated by the class  $\text{Alg}(\mathcal{S})$  and hence we have  $\text{Alg}(\mathcal{S}) \subseteq \mathbb{V}(\text{Alg}(\mathcal{S})) = \mathbf{K}_{\mathcal{S}}$ .*

**Definition 2.4** A sentential logic  $\mathcal{S}$  is said to be *fully selfextensional* (or *congruential*) if for every  $A \in \text{Alg}(\mathcal{S})$ , the Frege relation of the g-matrix  $\langle A, \text{Fi}_{\mathcal{S}}(A) \rangle$  is the identity relation.

## 2.2 Distributive nearlattices

Nearlattices and distributive nearlattices were studied by several authors [3, 8–13, 31, 32].

**Definition 2.5** [3] An algebra  $\langle A, m \rangle$  of type (3) is called a *nearlattice* if the following identities hold:

- (P1)  $m(x, y, x) = x$ ,
- (P2)  $m(m(x, y, z), m(y, m(u, x, z), z), w) = m(w, w, m(y, m(x, u, z), z))$ .

**Theorem 2.6** [13] *Let  $\langle A, m \rangle$  be an algebra of type (3) and let  $\vee$  be the binary operation on  $A$  defined by  $x \vee y := m(x, x, y)$ . Then,  $\langle A, m \rangle$  is a nearlattice if and only if  $\langle A, \vee \rangle$  is a join-semilattice where for every  $a \in A$ , the principal upset  $[a] = \{x \in A : a \leq x\}$  is a lattice with respect to the order  $\leq$  induced by  $\vee$ . Moreover, for all  $x, y, a \in A$ ,  $m(x, y, a) = (x \vee a) \wedge_a (y \vee a)$  where  $\wedge_a$  denotes the meet in  $[a]$ .*

Let  $\langle A, m \rangle$  be a nearlattice. We consider as the partial order of  $\langle A, m \rangle$  the order of the join-semilattice defined by  $m$ , that is,  $x \leq y$  if and only if  $y = x \vee y = m(x, x, y)$ . Moreover, for every element  $a \in A$  we have

$$x \wedge_a y = m(x, y, a)$$

for all  $x, y \in [a]$ . It should be noted that the meet  $x \wedge y$  exists in  $A$  if and only if  $\{x, y\}$  has a lower bound in  $A$ . Thus, the meet of  $x$  and  $y$  in  $[a]$  coincides with their meet in  $A$  for all  $x, y \in [a]$ , i.e.,  $x \wedge_a y = x \wedge y$ ; for instance, we have

$$m(a, b, c) = (a \vee c) \wedge_c (b \vee c) = (a \vee c) \wedge (b \vee c),$$


for all  $a, b, c \in A$ . This should be kept in mind since we will use it without mention.

**Definition 2.7** A nearlattice  $\langle A, m \rangle$  is said to be *distributive* if and only if it satisfies either of the following two equivalent identities:

- (P3)  $m(x, m(y, y, z), w) = m(m(x, y, w), m(x, y, w), m(x, z, w))$ ;
- (P4)  $m(x, x, m(y, z, w)) = m(m(x, x, y), m(x, x, z), w)$ .

Let us denote by  $\mathbb{DN}$  the variety of distributive nearlattices.

**Proposition 2.8** [13, Theorem 4] *A nearlattice  $\langle A, m \rangle$  is distributive if and only if for every  $a \in A$ , the lattice  $\langle [a], \wedge_a, \vee \rangle$  is distributive.*



**Fig. 1** A distributive nearlattice

*Example 2.9* In Fig. 1, it is shown a distributive nearlattice. For instance, we have

$$m(u, w, y) = (u \vee y) \wedge_y (w \vee y) = y \quad \text{and} \quad m(u, w, b) = (u \vee b) \wedge_b (w \vee b) = y.$$

That is,  $m(u, w, y) = u \wedge_y w = u \wedge w = u \wedge_b w = m(u, w, b)$ .

*Remark 2.10* Recall that a *Tarski algebra* (also called *implication algebra*) [1,2] can be defined as a binary algebra  $\langle A, \rightarrow \rangle$  satisfying some identities and, equivalently, it can be defined as a join-semilattice  $\langle A, \vee \rangle$  such that for every  $a \in A$ , the upset  $[a]$  is a Boolean algebra with respect to the order induced by  $\vee$ . Thus, we can noticed that the concept of distributive nearlattice is a natural generalisation of the notion of Tarski algebra.

**Definition 2.11** Let  $\langle A, m \rangle$  be a distributive nearlattice. A nonempty subset  $F \subseteq A$  is said to be a *filter* of  $A$  if (i)  $x \in F$  and  $x \leq y$  implies  $y \in F$ , and (ii) if  $x, y \in F$  and  $x \wedge y$  exists in  $A$ , then  $x \wedge y \in F$ .

Let us denote by  $\text{Fi}(A)$  the collection of all filters of a distributive nearlattice  $A$ . It is easy to check that for every distributive nearlattice  $A$  the intersection of any collection of filters is either a filter or an empty set. So, for every nonempty  $X \subseteq A$ , there exists the least filter containing  $X$ ; it is denoted by  $\text{Fi}_A(X)$ . If  $X = \{a_1, \dots, a_n\}$ , then we write  $\text{Fi}_A(a_1, \dots, a_n)$  instead of  $\text{Fi}_A(\{a_1, \dots, a_n\})$ ; moreover, it is easy to check that  $\text{Fi}_A(a) = [a]$ . There is a useful characterisation of the filter  $\text{Fi}_A(X)$  generated by  $X$ . Let  $X \subseteq A$  be nonempty. Then,

$$\text{Fi}_A(X) = \{a \in A : a = b_1 \wedge \dots \wedge b_k \text{ for some } b_1, \dots, b_k \in [X]\}.$$

where  $[X] = \{a \in A : a \geq x \text{ for some } x \in X\}$ , see [15].

**Proposition 2.12** [30] *Let  $A$  be a nearlattice and  $F \subseteq A$  be nonempty. Then, the following conditions are equivalent:*

1.  $F \in \text{Fi}(A)$ ;
2. if  $a, b \in F$ , then  $m(a, b, c) \in F$  for all  $c \in A$ .



Now we introduce the following definition that will be useful for what follows.

**Definition 2.13** Let  $\langle A, m \rangle$  be an algebra of type (3). For each integer  $n \geq 0$ , we define inductively, for all  $a_0, \dots, a_n, b \in A$ , an element  $m^n(a_0, \dots, a_n, b)$  as follows:

- $m^0(a_0, b) := m(a_0, a_0, b)$  and
- for  $n \geq 1$ ,  $m^n(a_0, \dots, a_n, b) := m(m^{n-1}(a_0, \dots, a_{n-1}, b), a_n, b)$ .

In particular, for a distributive nearlattice  $A$ , we get  $m^0(a_0, b) = a_0 \vee b$  and  $m^1(a_0, a_1, b) = m(a_0, a_1, b)$ . Indeed,

$$\begin{aligned} m^1(a_0, a_1, b) &= m(m^0(a_0, b), a_1, b) = m(a_0 \vee b, a_1, b) \\ &= (a_0 \vee b \vee b) \wedge_b (a_1 \vee b) = (a_0 \vee b) \wedge_b (a_1 \vee b) = m(a_0, a_1, b). \end{aligned}$$

The proofs of the following two propositions can be found in [31].

**Proposition 2.14** Let  $\langle A, m_A \rangle$  and  $\langle B, m_B \rangle$  be algebras of type (3) and  $h \in \text{Hom}(A, B)$ . Then, we have  $h(m_A^n(a_0, \dots, a_n, b)) = m_B^n(h(a_0), \dots, h(a_n), h(b))$  for all  $a_0, \dots, a_n, b \in A$ .

**Proposition 2.15** Let  $\langle A, m \rangle$  be a distributive nearlattice and  $a_0, \dots, a_n, a_{n+1}, a, b \in A$ . Then:

1.  $m^n(a_0, \dots, a_n, b) = (a_0 \vee b) \wedge_b \dots \wedge_b (a_n \vee b)$ ;
2.  $b \leq m^n(a_0, \dots, a_n, b)$ ;
3. if  $a \leq a_i$  for all  $i \in \{0, 1, \dots, n\}$ , then  $a \leq m^n(a_0, \dots, a_n, b)$ ;
4.  $m^{n+1}(a_0, \dots, a_{n+1}, b) \leq m^n(a_0, \dots, a_n, b)$ ;
5.  $m^n(a_0, \dots, a_n, b) = m^n(a_{\sigma(0)}, \dots, a_{\sigma(n)}, b)$ , for every permutation  $\sigma$  of  $\{0, 1, \dots, n\}$ ;
6. if  $a \in \text{Fi}_A(a_0, \dots, a_n)$ , then  $a \in \text{Fi}_A(m^n(a_0, \dots, a_n, a))$ ;
7.  $a \in \text{Fi}_A(a_0, \dots, a_n)$  if and only if  $a = m^n(a_0, \dots, a_n, a)$ .

### 3 Logics with a distributive nearlattice term

For a ternary term  $m$  of an algebraic language  $\mathcal{L}$ , we will consider the binary term  $\vee$  defined by  $x \vee y := m(x, x, y)$ . We also define, for every integer  $n \geq 0$  and variables  $x_0, \dots, x_n, x$ , the formula  $m^n(x_0, \dots, x_n, x)$  as follows:

- $m^0(x_0, x) := m(x_0, x_0, x)$
- for  $n \geq 1$ ,  $m^n(x_0, \dots, x_n, x) := m(m^{n-1}(x_0, \dots, x_{n-1}, x), x_n, x)$ .

**Definition 3.1** Let  $\mathcal{S}$  be a sentential logic over an algebraic language  $\mathcal{L}$ . A ternary term  $m$  of  $\mathcal{L}$  is said to be a *distributive nearlattice term* (DN-term) of  $\mathcal{S}$  if and only if  $\mathcal{S}$  satisfies the following properties:

- (A1)  $\varphi \vee \psi \vdash_{\mathcal{S}} \chi$  if and only if  $\varphi \vdash_{\mathcal{S}} \chi$  and  $\psi \vdash_{\mathcal{S}} \chi$ ;
- (A2)  $m(\varphi, \psi, \chi) \vdash_{\mathcal{S}} \varphi \vee \chi$  and  $m(\varphi, \psi, \chi) \vdash_{\mathcal{S}} \psi \vee \chi$ ;
- (A3)  $\varphi \vee \chi, \psi \vee \chi \vdash_{\mathcal{S}} m(\varphi, \psi, \chi)$ ;
- (A4) if  $\varphi_0, \dots, \varphi_n \vdash_{\mathcal{S}} \varphi$ , then  $m^n(\varphi_0, \dots, \varphi_n, \varphi) \vdash_{\mathcal{S}} \varphi$ ;

where  $x \vee y := m(x, x, y)$ .

Property (A1) is known in the literature as the *weak proof by cases property* [14, 40]. Moreover, (A1) implies that the following properties hold:  $\varphi \vee \varphi \vdash_{\mathcal{S}} \varphi$ ,  $\varphi \vdash_{\mathcal{S}} \varphi \vee \psi$ ,  $\varphi \vee \psi \vdash_{\mathcal{S}} \psi \vee \varphi$  and  $(\varphi \vee \psi) \vee \chi \vdash_{\mathcal{S}} \varphi \vee (\psi \vee \chi)$ . Then, the binary term  $\vee$  is a *disjunction* for the logic  $\mathcal{S}$ , and thus  $\mathcal{S}$  can be considered as a *disjunctive logic*, see [14, 17, 26, 40].

**Proposition 3.2** *If  $m$  is a DN-term of a sentential logic  $\mathcal{S}$ , then the following properties hold:*

1.  $m^n(\varphi_0, \dots, \varphi_n, \psi) \vdash_{\mathcal{S}} \varphi_i \vee \psi$ , for all  $i \in \{0, 1, \dots, n\}$ ;
2.  $\varphi_0 \vee \psi, \dots, \varphi_n \vee \psi \vdash_{\mathcal{S}} m^n(\varphi_0, \dots, \varphi_n, \psi)$ ;
3. if  $m^n(\varphi_0, \dots, \varphi_n, \varphi) \vdash_{\mathcal{S}} \varphi$ , then  $\varphi_0, \dots, \varphi_n \vdash_{\mathcal{S}} \varphi$ ;
4.  $\varphi \vdash_{\mathcal{S}} m^n(\varphi_0, \dots, \varphi_n, \varphi)$ .

*Proof* Properties (1) and (2) can be proved by induction on  $n$ . Property (1) follows from (A1) and (A2). Property (2) follows from (A1) and (A3). Properties (3) and (4) are consequences of (A1) and (2). □

**Proposition 3.3** *Let  $\mathcal{S}$  be a sentential logic. If  $m$  and  $m'$  are DN-terms of  $\mathcal{S}$ , then  $m(\varphi, \psi, \chi) \dashv\vdash_{\mathcal{S}} m'(\varphi, \psi, \chi)$ , for all  $\varphi, \psi, \chi \in Fm$ .*

*Proof* Let  $\varphi, \psi, \chi \in Fm$ . Since  $\varphi \vdash_{\mathcal{S}} \varphi \vee' \psi$  and  $\psi \vdash_{\mathcal{S}} \varphi \vee' \psi$ , it follows by (A1) that  $\varphi \vee \psi \vdash_{\mathcal{S}} \varphi \vee' \psi$ . Similarly, we have  $\varphi \vee' \psi \vdash_{\mathcal{S}} \varphi \vee \psi$ . Hence  $\varphi \vee \psi \dashv\vdash_{\mathcal{S}} \varphi \vee' \psi$ . Now, from this and using (A2) and (A3), we obtain

$$m(\varphi, \psi, \chi) \dashv\vdash_{\mathcal{S}} \{\varphi \vee \chi, \psi \vee \chi\} \dashv\vdash_{\mathcal{S}} \{\varphi \vee' \chi, \psi \vee' \chi\} \dashv\vdash_{\mathcal{S}} m'(\varphi, \psi, \chi). \quad \square$$

**Definition 3.4** A class of algebras  $\mathbf{K}$  of a given similarity type  $\mathcal{L}$  is called *distributive nearlattice-based* (DN-based for short) if there is a ternary term  $m$  of  $\mathcal{L}$  such that the distributive nearlattice equations (P1)–(P3) (Definitions 2.5 and 2.7) hold in  $\mathbf{K}$ . In this case, we will also say that  $\mathbf{K}$  is a *DN-class relative to  $m$*  and when there is not danger of confusion, we simply say that  $\mathbf{K}$  is a DN-class.

Notice that if  $\mathbf{K}$  is a DN-class relative to  $m$ , then for every algebra  $A \in \mathbf{K}$  the  $\{m\}$ -reduct  $\langle A, m^A \rangle$  is a distributive nearlattice. Moreover, we have that the variety  $\mathbb{V}(\mathbf{K})$  generated by a DN-class  $\mathbf{K}$  is also a DN-class.

**Definition 3.5** A sentential logic  $\mathcal{S}$  of type  $\mathcal{L}$  is said to be *distributive nearlattice-based* (DN-based for short) if there is a ternary term  $m$  and a DN-class  $\mathbf{K}$  of algebras of type  $\mathcal{L}$  such that the following condition holds for every  $n \geq 0$  and for all formulas  $\varphi_0, \dots, \varphi_n, \varphi$ :

$$\varphi_0, \dots, \varphi_n \vdash_{\mathcal{S}} \varphi \iff (\forall A \in \mathbf{K})(\forall h \in \text{Hom}(Fm, A)) \quad (3.1)$$

$$m^n(h(\varphi_0), \dots, h(\varphi_n), h(\varphi)) \leq h(\varphi).$$

We will say that  $\mathcal{S}$  is *DN-based relative to  $m$  and  $\mathbf{K}$* .

It should be noted, by property (5) of Proposition 2.15, that condition (3.1) is independent of the order in which the formulas  $\varphi_0, \dots, \varphi_n$  are taken.

Let  $\mathcal{S}$  be a DN-based logic relative to  $m$  and  $\mathbf{K}$ . It is easy to check that for every formulas  $\varphi$  and  $\psi$ ,

$$\varphi \vdash_{\mathcal{S}} \psi \iff (\forall A \in \mathbf{K})(\forall h \in \text{Hom}(Fm, A))h(\varphi) \leq h(\psi).$$

Then, we obtain that for all  $\varphi, \psi \in Fm$

$$\varphi \dashv\vdash_{\mathcal{S}} \psi \iff \mathbf{K} \models \varphi \approx \psi \iff \mathbb{V}(\mathbf{K}) \models \varphi \approx \psi. \tag{3.2}$$

Noticed that (3.2) is independent of the term  $m$ .

From (3.1) and by property (2) of Proposition 2.15, we have

$$\begin{aligned} \varphi_0, \dots, \varphi_n \vdash_{\mathcal{S}} \varphi &\iff (\forall A \in \mathbf{K})(\forall h \in \text{Hom}(Fm, A)) \\ &\quad m^n(h(\varphi_0), \dots, h(\varphi_n), h(\varphi)) = h(\varphi) \\ &\iff \mathbf{K} \models m^n(\varphi_0, \dots, \varphi_n, \varphi) \approx \varphi \\ &\iff \mathbb{V}(\mathbf{K}) \models m^n(\varphi_0, \dots, \varphi_n, \varphi) \approx \varphi. \end{aligned}$$

Hence,  $\mathcal{S}$  is also DN-based relative to the variety  $\mathbb{V}(\mathbf{K})$  generated by  $\mathbf{K}$ . Moreover, by (3.2), we can see that the variety to which  $\mathcal{S}$  is DN-based is unique. So, let us denote the only variety relative to which  $\mathcal{S}$  is DN-based by  $\mathbb{V}(\mathcal{S})$ .

**Proposition 3.6** *Let  $\mathcal{S}$  be a DN-based logic relative to  $m$ . Then,  $m$  is a DN-term of  $\mathcal{S}$ .*

*Proof* We have that  $\mathcal{S}$  is DN-based relative to  $\mathbb{V}(\mathcal{S})$  and the ternary term  $m$ . Property (A1) is a consequence of the fact that for every  $A \in \mathbb{V}(\mathcal{S})$ , the  $\{\vee\}$ -reduct  $\langle A, \vee^A \rangle$  is a join-semilattice. Property (A2) holds because for every  $A \in \mathbb{V}(\mathcal{S})$  and all  $a, b, c \in A$ , we have  $m^A(a, b, c) = (a \vee c) \wedge (b \vee c) \leq a \vee c, b \vee c$ . In order to prove (A3), let  $\varphi, \psi, \chi \in Fm$ . Let  $A \in \mathbb{V}(\mathcal{S})$  and  $h \in \text{Hom}(Fm, A)$ . Assume that  $h(\varphi) = a, h(\psi) = b$  and  $h(\chi) = c$ . So, we need to show that  $m(a \vee c, b \vee c, m(a, b, c)) \leq m(a, b, c)$ . Now, by condition (P4), we have

$$\begin{aligned} m(a \vee c, b \vee c, m(a, b, c)) &= c \vee m(a, b, m(a, b, c)) = m(a, b, m(a, b, c)) \\ &= (a \vee m(a, b, c)) \wedge (b \vee m(a, b, c)) = (a \vee c) \wedge (b \vee c) = m(a, b, c). \end{aligned}$$

Hence property (A3) holds. Lastly, property (A4) is an immediate consequence by (3.1). □

By the previous proposition and from (3.2), we obtain that if  $\mathcal{S}$  is a DN-based logic relative to ternary terms  $m$  and  $m'$ , then for every  $A \in \mathbb{V}(\mathcal{S})$ , we have that the ternary operations  $m^A$  and  $m'^A$  coincide. Hence, we can say simply that a logic  $\mathcal{S}$  is DN-based.

**Proposition 3.7** *Let  $\mathcal{S}$  be a DN-based logic. Then  $\mathcal{S}$  is selfextensional and  $\mathbb{V}(\mathcal{S}) = \mathbf{K}_{\mathcal{S}}$ .*

*Proof* By definition of the Frege relation and from (3.2), we have  $(\varphi, \psi) \in \Lambda(\mathcal{S}) \iff \mathbb{V}(\mathcal{S}) \models \varphi \approx \psi$ . Hence, we obtain that  $\Lambda(\mathcal{S})$  is a congruence on  $Fm$ . Therefore  $\mathcal{S}$  is selfextensional. Now, since  $\mathcal{S}$  selfextensional, it follows that  $\varphi \dashv\vdash_{\mathcal{S}} \psi \iff K_{\mathcal{S}} \models \varphi \approx \psi$ . Then, by (3.2) again, we obtain that  $\mathbb{V}(\mathcal{S}) \models \varphi \approx \psi \iff \varphi \dashv\vdash_{\mathcal{S}} \psi \iff K_{\mathcal{S}} \models \varphi \approx \psi$ . Therefore  $\mathbb{V}(\mathcal{S}) = K_{\mathcal{S}}$ .  $\square$

Now we are ready to show one of the main results of this paper.

**Theorem 3.8** *Let  $\mathcal{S}$  be a sentential logic. Then,  $\mathcal{S}$  is a selfextensional logic with a DN-term  $m$  if and only if  $\mathcal{S}$  is a DN-based logic relative to  $m$ .*

*Proof* The implication from right to left is a consequence from Propositions 3.6 and 3.7. Now we assume that  $\mathcal{S}$  is selfextensional and  $m$  is a DN-term of  $\mathcal{S}$ . First, since  $\mathcal{S}$  is selfextensional, it follows that  $\Lambda(\mathcal{S})$  is a congruence on  $Fm$  and hence, we can consider the quotient algebra  $Fm^* := Fm/\Lambda(\mathcal{S})$ . Let us show that  $\langle Fm^*, m^* \rangle$ , with  $m^*(\overline{\varphi}, \overline{\psi}, \overline{\chi}) := \overline{m(\varphi, \psi, \chi)}$  ( $\overline{\varphi}$  denotes the equivalent class of  $\varphi$  in  $Fm/\Lambda(\mathcal{S})$ ), is a distributive nearlattice. By (A1),  $\mathcal{S}$  satisfies the following properties:  $\varphi \vee \varphi \vdash_{\mathcal{S}} \varphi$ ,  $\varphi \vdash_{\mathcal{S}} \varphi \vee \psi$ ,  $\varphi \vee \psi \vdash_{\mathcal{S}} \psi \vee \varphi$  and  $\varphi \vee (\psi \vee \chi) \dashv\vdash_{\mathcal{S}} (\varphi \vee \psi) \vee \chi$ . Thus, it is easy to check that  $\langle Fm^*, \vee^* \rangle$ , with  $\overline{\varphi} \vee^* \overline{\psi} := m^*(\overline{\varphi}, \overline{\varphi}, \overline{\psi})$ , is a join-semilattice. Let us denote by  $\leq$  the order of  $\langle Fm^*, \vee^* \rangle$ . Let  $\chi \in Fm$ . We prove that  $[\overline{\chi}] = \{\overline{\varphi} \in Fm^* : \overline{\chi} \leq \overline{\varphi}\} = \{\overline{\varphi} \in Fm^* : \chi \vdash_{\mathcal{S}} \varphi\}$  is a distributive lattice. In order to prove that  $[\overline{\chi}]$  is a lattice, we need only to show that there exists the meet in  $[\overline{\chi}]$ . Let  $\overline{\varphi}, \overline{\psi} \in [\overline{\chi}]$ . So  $\chi \vdash_{\mathcal{S}} \varphi, \psi$ . Let us prove that  $m^*(\overline{\varphi}, \overline{\psi}, \overline{\chi})$  is the meet of  $\overline{\varphi}$  and  $\overline{\psi}$  in  $[\overline{\chi}]$ . By (A1) and (A2) we have  $m(\varphi, \overline{\psi}, \chi) \vdash_{\mathcal{S}} \varphi \vee \chi \vdash_{\mathcal{S}} \varphi$  and  $m(\varphi, \psi, \chi) \vdash_{\mathcal{S}} \psi \vee \chi \vdash_{\mathcal{S}} \psi$ . Thus  $m^*(\overline{\varphi}, \overline{\psi}, \overline{\chi}) \leq \overline{\varphi}, \overline{\psi}$ . Let  $\overline{\gamma} \in [\overline{\chi}]$  be such that  $\overline{\gamma} \leq \overline{\varphi}, \overline{\psi}$ . So  $\chi \vdash_{\mathcal{S}} \gamma$  and  $\gamma \vdash_{\mathcal{S}} \varphi, \psi$ . Then  $\gamma \vdash_{\mathcal{S}} \varphi \vee \chi, \psi \vee \chi$ . By (A3) we obtain that  $\gamma \vdash_{\mathcal{S}} m(\varphi, \psi, \chi)$ , that is,  $\overline{\gamma} \leq m^*(\overline{\varphi}, \overline{\psi}, \overline{\chi})$ . Hence  $m^*(\overline{\varphi}, \overline{\psi}, \overline{\chi}) = \overline{\varphi} \wedge_{[\overline{\chi}]} \overline{\psi}$ . Then, by Theorem 2.6, we conclude that  $\langle Fm^*, m^* \rangle$  is a nearlattice. Now we show that condition (P4) holds in  $\langle Fm^*, m^* \rangle$ . Let  $\varphi, \psi, \gamma, \chi \in Fm$ . Since  $\langle Fm^*, m^* \rangle$  is a nearlattice, it follows that  $\overline{\varphi} \vee^* m^*(\overline{\psi}, \overline{\gamma}, \overline{\chi}) \leq m^*(\overline{\varphi} \vee^* \overline{\psi}, \overline{\varphi} \vee^* \overline{\gamma}, \overline{\chi})$ . In order to prove the inverse inequality, we need to show that  $m(\varphi \vee \psi, \varphi \vee \gamma, \chi) \vdash_{\mathcal{S}} \varphi \vee m(\psi, \gamma, \chi)$ . By (A1), we have  $\psi, \gamma \vdash_{\mathcal{S}} \psi \vee \chi, \gamma \vee \chi$ , and from (A3) we obtain that  $\psi, \gamma \vdash_{\mathcal{S}} m(\psi, \gamma, \chi)$ . Thus  $\psi, \gamma \vdash_{\mathcal{S}} \varphi \vee m(\psi, \gamma, \chi)$ . By (A4), it follows that

$$m(\psi, \gamma, \varphi \vee m(\psi, \gamma, \chi)) \vdash_{\mathcal{S}} \varphi \vee m(\psi, \gamma, \chi). \tag{3.3}$$

By (A1) and (A3), we can deduce  $\varphi \vee \psi \vee \chi \vdash_{\mathcal{S}} \varphi \vee \psi \vee m(\psi, \gamma, \chi)$  and  $\varphi \vee \gamma \vee \chi \vdash_{\mathcal{S}} \varphi \vee \gamma \vee m(\psi, \gamma, \chi)$ . Then, by (A1)–(A3) and (3.3), we have

$$\begin{aligned} m(\varphi \vee \psi, \varphi \vee \gamma, \chi) &\vdash_{\mathcal{S}} \varphi \vee \psi \vee \chi, \varphi \vee \gamma \vee \chi \\ &\vdash_{\mathcal{S}} \varphi \vee \psi \vee m(\psi, \gamma, \chi), \varphi \vee \gamma \vee m(\psi, \gamma, \chi) \\ &\vdash_{\mathcal{S}} \psi \vee \varphi \vee m(\psi, \gamma, \chi), \gamma \vee \varphi \vee m(\psi, \gamma, \chi) \\ &\vdash_{\mathcal{S}} m(\psi, \gamma, \varphi \vee m(\psi, \gamma, \chi)) \\ &\vdash_{\mathcal{S}} \varphi \vee m(\psi, \gamma, \chi). \end{aligned}$$

Hence, we have proved that  $\langle Fm^*, m^* \rangle$  is a distributive nearlattice. Finally, we prove that  $\mathcal{S}$  is DN-based relative to  $\{Fm^*\}$  and  $m$ . Let  $\varphi_0, \dots, \varphi_n, \varphi \in Fm$ . From property

(A4), (3) of Proposition 3.2, (2) of Proposition 2.15 and since  $\mathcal{S}$  is selfextensional, it follows that

$$\begin{aligned} \varphi_0, \dots, \varphi_n \vdash_{\mathcal{S}} \varphi &\iff m^n(\varphi_0, \dots, \varphi_n, \varphi) \vdash_{\mathcal{S}} \varphi \\ &\iff \overline{m^n(\varphi_0, \dots, \varphi_n, \varphi)} \leq \overline{\varphi} \iff \overline{m^n(\varphi_0, \dots, \varphi_n, \varphi)} = \overline{\varphi} \\ &\iff Fm^* \models m^n(\varphi_0, \dots, \varphi_n, \varphi) \approx \varphi \\ &\iff (\forall h \in \text{Hom}(Fm, Fm^*))(m^n(h\varphi_0, \dots, h\varphi_n, h\varphi) = h\varphi) \\ &\iff (\forall h \in \text{Hom}(Fm, Fm^*))(m^n(h\varphi_0, \dots, h\varphi_n, h\varphi) \leq h\varphi). \end{aligned}$$

This completes the proof. □

Our next aim is to prove that every selfextensional logic  $\mathcal{S}$  with a DN-term is fully selfextensional and the class  $\text{Alg}(\mathcal{S})$  is a variety. Notice, by the previous theorem and Proposition 3.7, that for every selfextensional logic  $\mathcal{S}$  with a DN-term  $m$  the  $m$ -reducts of the algebras of its intrinsic variety  $\mathbf{K}_{\mathcal{S}}$  are distributive nearlattices and  $\mathcal{S}$  is DN-based relative to  $\mathbf{K}_{\mathcal{S}}$ .

**Proposition 3.9** *Let  $\mathcal{S}$  be a DN-based logic relative to  $m$ . Then, for every algebra  $A \in \mathbf{K}_{\mathcal{S}}$ , the nonempty  $\mathcal{S}$ -filters of  $A$  are exactly the filters of the  $\{m\}$ -reduct distributive nearlattice  $\langle A, m^A \rangle$ , i.e.,  $\text{Fi}_{\mathcal{S}}(A) \setminus \{\emptyset\} = \text{Fi}(A)$ .*

*Proof* Let  $A \in \mathbf{K}_{\mathcal{S}}$ . Let  $F \in \text{Fi}(A)$ . Let  $\varphi_0, \dots, \varphi_n, \varphi \in Fm$  be such that  $\varphi_0, \dots, \varphi_n \vdash_{\mathcal{S}} \varphi$  and let  $h \in \text{Hom}(Fm, A)$  be such that  $h(\varphi_i) \in F$  for all  $i = 0, 1, \dots, n$ . By (3.1), we have  $m^n(h(\varphi_0), \dots, h(\varphi_n), h(\varphi)) \leq h(\varphi)$ . Since  $F$  is a filter of the nearlattice  $A$  and  $h(\varphi_0), \dots, h(\varphi_n) \in F$ , it follows by Proposition 2.12 that  $m^n(h(\varphi_0), \dots, h(\varphi_n), h(\varphi)) \in F$ . Then  $h(\varphi) \in F$ . Hence  $F \in \text{Fi}_{\mathcal{S}}(A)$ . Conversely, let now  $F \in \text{Fi}_{\mathcal{S}}(A)$  be nonempty. Let  $a, b \in F$  and  $c \in A$ . By (A1) and (A3) we have, for variables  $x, y$  and  $z$ , that  $x, y \vdash_{\mathcal{S}} \{x \vee z, y \vee z\} \vdash_{\mathcal{S}} m(x, y, z)$ . By taking  $h \in \text{Hom}(Fm, A)$  such that  $h(x) = a, h(y) = b$  and  $h(z) = c$ , we obtain that  $h(x), h(y) \in F$  and hence  $h(m(x, y, z)) \in F$ , i.e.,  $m(a, b, c) \in F$ . Therefore,  $F \in \text{Fi}(A)$ . □

**Theorem 3.10** *Let  $\mathcal{S}$  be a DN-based logic. Then:*

1.  $\text{Alg}(\mathcal{S}) = \mathbf{K}_{\mathcal{S}}$ ;
2.  $\text{Alg}(\mathcal{S})$  is a variety;
3.  $\mathcal{S}$  is DN-based relative to  $\text{Alg}(\mathcal{S})$ .

*Proof* (1) We know by Lemma 2.3 that  $\text{Alg}(\mathcal{S}) \subseteq \mathbf{K}_{\mathcal{S}}$ . Let  $A \in \mathbf{K}_{\mathcal{S}}$ . From Proposition 3.9 we can easily deduce that the  $g$ -matrix  $\langle A, \text{Fi}_{\mathcal{S}}(A) \rangle$  is a reduced  $g$ -model of  $\mathcal{S}$ . Hence  $A \in \text{Alg}(\mathcal{S})$ . Properties (2) and (3) are immediate consequences of (1). □

**Corollary 3.11** *If  $\mathcal{S}$  is a DN-based logic, then  $\mathcal{S}$  is fully selfextensional.*

*Proof* Recall the definition of fully selfextensionality, see Definition 2.4. Let  $A \in \text{Alg}(\mathcal{S})$ . So  $A \in \mathbf{K}_{\mathcal{S}}$ . Then, by Proposition 3.9, it is easy check that  $\Lambda_A(\text{Fi}_{\mathcal{S}}(A)) = \text{Id}_A$ . Hence  $\mathcal{S}$  is fully-selfextensional. □

We have characterised selfextensional logics with a DN-term as those logics that are DN-based concerning their canonical class of algebras. As happens in the setting of selfextensional logics with a conjunction [34], two different sentential logics  $\mathcal{S}$  and  $\mathcal{S}'$  can be DN-based relative to the same DN-based variety  $\mathbf{K}$ . The unique possible case for this is when one of them has theorems and the other has not. Now we will see under what conditions the uniqueness can be obtained.

Let  $\mathbf{K}$  be a DN-based variety relative to a ternary term  $m$ . Let us define the sentential logic  $\mathcal{S}_{\mathbf{K}} = \langle Fm, \vdash_{\mathbf{K}} \rangle$  as follows: let  $\varphi_0, \dots, \varphi_n, \varphi \in Fm$ ,

$$\varphi_0, \dots, \varphi_n \vdash_{\mathbf{K}} \varphi \iff (\forall A \in \mathbf{K})(\forall h \in \text{Hom}(Fm, A)) \quad (3.4)$$

$$m^n(h\varphi_0, \dots, h\varphi_n, h\varphi) \leq h\varphi$$

and

$$\emptyset \vdash_{\mathbf{K}} \varphi \iff (\forall A \in \mathbf{K})(\forall h \in \text{Hom}(Fm, A)) \quad (3.5)$$

$$(\forall a \in A)(a \leq h\varphi).$$

Now, for every  $\Gamma \subseteq Fm$ ,  $\Gamma \vdash_{\mathbf{K}} \varphi$  if and only if there is a finite  $\Gamma_0 \subseteq \Gamma$  such that  $\Gamma_0 \vdash_{\mathbf{K}} \varphi$ . Notice that if  $\mathcal{S}_{\mathbf{K}}$  has a theorem, then for every algebra  $A \in \mathbf{K}$  the  $\{m\}$ -reduct nearlattice  $\langle A, m^A \rangle$  has a greatest element. Moreover, since  $\mathbf{K}$  is a variety, it follows that  $\mathbf{K}$  is the unique DN-based variety to which  $\mathcal{S}_{\mathbf{K}}$  is DN-based and hence, by Theorem 3.10, we have  $\mathbf{K}_{\mathcal{S}_{\mathbf{K}}} = \text{Alg}(\mathcal{S}_{\mathbf{K}}) = \mathbf{K}$ .

A sentential logic  $\mathcal{S}$  is said to be *non-pseudo axiomatic* [36] if for every formula  $\varphi$ ,  $\varphi$  is a theorem if and only if  $\varphi$  is derivable from every formula  $(\psi \vdash_{\mathcal{S}} \varphi$  for all formulas  $\psi)$ , or equivalently if the intersection of all its nonempty theories is the set of theorems. Notice that every sentential logic with theorems is non-pseudo axiomatic. The following proposition is an immediate consequence from (3.4) and (3.5), and thus we omit its proof.

**Proposition 3.12** *Let  $\mathcal{L}$  be an algebraic language and  $m$  a ternary term of  $\mathcal{L}$ . If  $\mathbf{K}$  is a DN-based variety relative to  $m$ , then  $\mathcal{S}_{\mathbf{K}}$  is the unique non-pseudo axiomatic sentential logic which is DN-based relative to  $\mathbf{K}$  and  $m$ ; moreover  $\mathbf{K}_{\mathcal{S}_{\mathbf{K}}} = \mathbf{K}$ . If  $\mathcal{S}$  is a DN-based and non-pseudo axiomatic logic relative to  $m$ , then  $\mathcal{S}_{\mathbf{K}_{\mathcal{S}}} = \mathcal{S}$ .*

Hence, under the condition of being non-pseudo axiomatic, we obtain the following kind of uniqueness for DN-based logics: different non-pseudo axiomatic logics must be DN-based relative to different DN-based varieties.

Now, we show a bijective correspondence between the class of DN-based and non-pseudo axiomatic logics and the class of subvarieties of the variety axiomatized by the equations (P1)-(P3).

A sentential logic  $\mathcal{S}'$  is said to be an *extension* of a sentential logic  $\mathcal{S}$  if and only if for every  $\Gamma \cup \{\varphi\} \subseteq Fm$ ,  $\Gamma \vdash_{\mathcal{S}} \varphi$  implies  $\Gamma \vdash_{\mathcal{S}'} \varphi$ .

**Lemma 3.13** *Let  $\mathcal{S}$  and  $\mathcal{S}'$  be DN-based and non-pseudo axiomatic logics. Then,  $\Lambda(\mathcal{S}) \subseteq \Lambda(\mathcal{S}')$  if and only if  $\mathcal{S}'$  is an extension of  $\mathcal{S}$ .*

*Proof* It is immediate that if  $S'$  is an extension of  $S$ , then  $\Lambda(S) \subseteq \Lambda(S')$ . So, we need to prove the implication from left to right. Assume that  $\Lambda(S) \subseteq \Lambda(S')$ . By property (A4) and properties (3) and (4) of Proposition 3.2, it follows that

$$\begin{aligned} \varphi_0, \dots, \varphi_n \vdash_S \varphi &\iff m^n(\varphi_0, \dots, \varphi_n, \varphi) \vdash_S \varphi \iff \\ &m^n(\varphi_0, \dots, \varphi_n, \varphi) \dashv\vdash_S \varphi \implies m^n(\varphi_0, \dots, \varphi_n, \varphi) \dashv\vdash_{S'} \varphi \iff \\ &m^n(\varphi_0, \dots, \varphi_n, \varphi) \vdash_{S'} \varphi \iff \varphi_0, \dots, \varphi_n \vdash_{S'} \varphi. \end{aligned}$$

Now, if  $\emptyset \vdash_S \varphi$ , then  $\psi \vdash_S \varphi$  for every formula  $\psi$ . By the above, we obtain that  $\psi \vdash_{S'} \varphi$  for every formula  $\psi$ . Now, since  $S'$  is non-pseudo axiomatic, it follows that  $\emptyset \vdash_{S'} \varphi$ . Hence, we have proved that  $S'$  is an extension of  $S$ .  $\square$

Let  $\mathcal{L}$  be an algebraic language and  $m$  a ternary term of  $\mathcal{L}$ . We set

- $\mathbb{S}_m(\mathcal{L}) := \{S : S \text{ is a non-pseudo-axiomatic logic over } \mathcal{L} \text{ and DN-based relative to } m\}$
- $\mathbb{K}_m(\mathcal{L}) := \{K : K \text{ is a subvariety of the variety over } \mathcal{L} \text{ axiomatized by the equations (P1)-(P3) with regard to } m\}$ .

We consider  $\mathbb{S}_m(\mathcal{L})$  ordered by the extension order, i.e.,  $S \leq S'$  if and only if  $S'$  is an extension of  $S$  and  $\mathbb{K}_m(\mathcal{L})$  ordered by the inclusion order. Now, we are in a position to establish and prove the announced result above.

**Theorem 3.14** *Let  $\mathcal{L}$  be an algebraic language and  $m$  a ternary term of  $\mathcal{L}$ . Then, the map  $F: \mathbb{S}_m(\mathcal{L}) \rightarrow \mathbb{K}_m(\mathcal{L})$  defined by  $F(S) = K_S$ , is a dual order isomorphism.*

*Proof* By Proposition 3.7, we have that  $F$  is well defined, and by Proposition 3.12 we obtain that  $F$  is an onto map. Let  $S, S' \in \mathbb{S}_m(\mathcal{L})$ . Then, by Lemma 3.13 and using that  $S$  and  $S'$  are selfextensional, we have

$$\begin{aligned} S \leq S' &\iff \Lambda(S) \subseteq \Lambda(S') \\ &\iff (\forall \varphi, \psi \in Fm)(K_S \models \varphi \approx \psi \implies K_{S'} \models \varphi \approx \psi) \\ &\iff K_{S'} \subseteq K_S. \end{aligned}$$

Therefore,  $F$  is a dual order isomorphism.  $\square$

## 4 Two examples

### 4.1 The logic of distributive nearlattices

In [31], it is defined a sentential logic  $S_{dn}$  through a Gentzen calculus, which can be considered as naturally associated with the variety of distributive nearlattices. There, the logic  $S_{dn}$  is denoted by  $S_{\mathbb{DN}}$ , but here  $S_{\mathbb{DN}}$  has a specific definition, see (3.4) and (3.5). Let us show that  $S_{dn}$  is the unique non-pseudo axiomatic DN-based logic relative to the variety of distributive nearlattices  $\mathbb{DN}$ . To this end, we need to introduce some basic notions of Gentzen calculus; we refer the reader to [26,31] for more information.

Let  $Fm$  be the algebra of formulas of a given algebraic similarity type  $\mathcal{L}$ . For our purpose, we will consider a *sequent* of type  $\mathcal{L}$  to be a pair  $\langle \Gamma, \varphi \rangle$  where  $\Gamma$  is a (possible empty) finite set of formulas and  $\varphi$  is a formula. As usual, we write  $\Gamma \triangleright \varphi$  instead of  $\langle \Gamma, \varphi \rangle$ . Let us denote by  $\text{Seq}(\mathcal{L})$  the collection of all sequents. A *Gentzen-style rule* is a pair  $\langle X, \Gamma \triangleright \varphi \rangle$  where  $X$  is a (possible empty) finite set of sequents and  $\Gamma \triangleright \varphi$  is a sequent. As usual, we shall use the standard fraction notation for Gentzen-style rules:

$$\frac{\Gamma_0 \triangleright \varphi_0, \dots, \Gamma_{n-1} \triangleright \varphi_{n-1}}{\Gamma \triangleright \varphi} \tag{4.1}$$

A *substitution instance of a Gentzen-style rule*  $\langle X, \Gamma \triangleright \varphi \rangle$  is a Gentzen-style rule of the form  $\langle \sigma[X], \sigma[\Gamma] \triangleright \sigma(\varphi) \rangle$  for some substitution  $\sigma \in \text{Hom}(Fm, Fm)$ . A *Gentzen calculus* is a set of Gentzen-style rules. Given a Gentzen calculus  $\mathbf{G}$ , the notion of a formal proof can be defined as usual. That is, a *pf* in the Gentzen calculus  $\mathbf{G}$  from a set of sequents  $X$  is a finite sequence of sequents each one of whose elements is a substitution instance of a rule of  $\mathbf{G}$  or a sequent in  $X$  or is obtained by applying a substitution instance of a rule of  $\mathbf{G}$  to previous elements in the sequence. A sequent  $\Gamma \triangleright \varphi$  is *derivable in  $\mathbf{G}$*  from a set of sequents  $X$  if there is a proof in  $\mathbf{G}$  from  $X$  whose last sequent in the proof is  $\Gamma \triangleright \varphi$ . We express this writing  $X \vdash_{\mathbf{G}} \Gamma \triangleright \varphi$ .

**Definition 4.1** A *Gentzen system* is a pair  $\mathcal{G} = \langle Fm, \vdash_{\mathcal{G}} \rangle$  where  $\vdash_{\mathcal{G}}$  is a finitary and structural (substitution-invariant) consequence relation on the set  $\text{Seq}(\mathcal{L})$  which in addition satisfies the following *structural rules*: for every  $\Gamma \cup \{\varphi, \psi\} \subseteq Fm$ ,

$$\text{(Axiom)} \frac{\emptyset}{\varphi \triangleright \varphi} \quad \text{(Weakening)} \frac{\Gamma \triangleright \varphi}{\Gamma, \psi \triangleright \varphi} \quad \text{(Cut)} \frac{\Gamma \triangleright \varphi \quad \Gamma, \varphi \triangleright \psi}{\Gamma \triangleright \psi}$$

We say that a Gentzen system  $\mathcal{G} = \langle Fm, \vdash_{\mathcal{G}} \rangle$  *satisfies a Gentzen-style rule* of type (4.1) or that (4.1) is a *Gentzen-style rule of  $\mathcal{G}$*  if  $\Gamma_0 \triangleright \varphi_0, \dots, \Gamma_{n-1} \triangleright \varphi_{n-1} \vdash_{\mathcal{G}} \Gamma \triangleright \varphi$  and we say that a sequent  $\Gamma \triangleright \varphi$  is a *derivable sequent* of  $\mathcal{G}$  when  $\emptyset \vdash_{\mathcal{G}} \Gamma \triangleright \varphi$ .

Let  $\mathbf{G}$  be a Gentzen calculus with the structural rules of (Axiom), (Weakening) and (Cut). Hence,  $\mathbf{G}$  defines in a standard way the Gentzen system  $\mathcal{G}_{\mathbf{G}} = \langle Fm, \vdash_{\mathbf{G}} \rangle$  (see [26, 39]).

Now let  $\mathcal{L} = \{m\}$  with  $m$  a ternary connective.

**Definition 4.2** [31, Definition 4.2] Let  $\mathcal{G}_{\text{DN}} = \langle Fm, \vdash_{\text{DN}} \rangle$  be the Gentzen system defined by the following Gentzen-style rules: the structural rules (Axiom), (Weakening) and (Cut) and the following rules

$$\begin{aligned} (\vee \triangleright) & \frac{\varphi \triangleright \chi \quad \psi \triangleright \chi}{\varphi \vee \psi \triangleright \chi} & (\triangleright \vee) & \frac{\Gamma \triangleright \varphi \quad \Gamma \triangleright \varphi}{\Gamma \triangleright \varphi \vee \psi \quad \Gamma \triangleright \varphi \vee \psi} \\ (m \triangleright) & \frac{}{m(\varphi, \psi, \chi) \triangleright \varphi \vee \chi \quad m(\varphi, \psi, \chi) \triangleright \psi \vee \chi} \\ (\triangleright m) & \frac{\Gamma \triangleright \varphi \vee \chi \quad \Gamma \triangleright \psi \vee \chi}{\Gamma \triangleright m(\varphi, \psi, \chi)} & (m^n \triangleright) & \frac{\varphi_1, \dots, \varphi_n \triangleright \varphi}{m^n(\varphi_0, \dots, \varphi_n, \varphi) \triangleright \varphi} \end{aligned}$$



Now, the sentential logic  $\mathcal{S}_{dn} = \langle Fm, \vdash_{dn} \rangle$  is defined as follows: for all  $\Gamma \cup \{\varphi\} \subseteq Fm$ ,

$$\Gamma \vdash_{dn} \varphi \iff \text{there is a finite } \Gamma_0 \subseteq \Gamma \text{ such that } \vdash_{\mathbb{DN}} \Gamma_0 \triangleright \varphi.$$

It follows directly that the connective  $m$  is a DN-term of  $\mathcal{S}_{dn}$ . Indeed, property (A1) follows from rules  $(\vee \triangleright)$  and  $(\triangleright \vee)$ ; (A2) follows from  $(m \triangleright)$ ; property (A3) is a consequence of  $(\triangleright m)$ ; and lastly, (A4) follows from the rule  $(m^n \triangleright)$ .

**Theorem 4.3** [31, Theorem 4.15] *The sentential logic  $\mathcal{S}_{dn}$  has the following properties:*

1.  $\text{Alg}(\mathcal{S}_{dn}) = \mathbb{DN}$ ;
2. for all  $\varphi_0, \dots, \varphi_n, \varphi \in Fm$ ,

$$\varphi_0, \dots, \varphi_n \vdash_{dn} \varphi \iff \mathbb{DN} \models m^n(\varphi_0, \dots, \varphi_n, \varphi) \approx \varphi.$$

3. for every  $A \in \text{Alg}(\mathcal{S}_{dn})$ ,  $\text{Fi}_{\mathcal{S}_{dn}}(A) = \text{Fi}(A) \cup \{\emptyset\}$ ;
4.  $\mathcal{S}_{dn}$  is fully selfextensional.

Therefore, by condition (2) of the previous theorem, we have that the logic  $\mathcal{S}_{dn}$  is DN-based relative to the variety  $\mathbb{DN}$ . Moreover, since  $\mathcal{S}_{dn}$  is non-pseudo axiomatic, it follows by Proposition 3.12 that  $\mathcal{S}_{dn} = \mathcal{S}_{\mathbb{DN}}$ .

### 4.2 Modal distributive nearlattices

In [7] the author introduces a notion of a necessity modal operator defined on a distributive nearlattice, and he studies these structures through a topological duality for the category of distributive nearlattices.

Let us consider the algebraic language  $\mathcal{L} = \{m, \Box, \top\}$  of type  $(3,1,0)$ .

**Definition 4.4** [7] An algebra  $\langle A, m, \Box, 1 \rangle$  is said to be a  $\Box$ -modal distributive nearlattice if  $\langle A, m, 1 \rangle$  is a distributive nearlattice with a greatest element 1, and the following conditions hold:

1.  $\Box 1 = 1$ ;
2. for all  $a, b \in A$  such that  $a \wedge b$  exists,  $\Box(a \wedge b) = \Box a \wedge \Box b$ .

We denote by  $\Box\mathbb{DN}$  the collection of all  $\Box$ -modal distributive nearlattices. Let us show that  $\Box\mathbb{DN}$  is a variety.

**Proposition 4.5** *Let  $\langle A, m, \Box, 1 \rangle$  be an algebra such that  $\langle A, m, 1 \rangle$  is a distributive nearlattice with a greatest element 1 and  $\Box 1 = 1$ . Then,  $\langle A, m, \Box, 1 \rangle \in \Box\mathbb{DN}$  if and only if the following identity holds in  $\langle A, m, \Box, 1 \rangle$ :*

$$\Box m(x, y, z) = m(\Box(x \vee z), \Box(y \vee z), \Box z). \tag{M}$$

Therefore,  $\Box\mathbb{DN}$  is the variety defined by identities of distributive nearlattices with a greatest element, and the identities  $\Box 1 = 1$  and (M).

*Proof* First assume that  $\langle A, m, \Box, 1 \rangle \in \Box\mathbb{DN}$ . Let  $a, b, c \in A$ . Since the operator  $\Box$  is order-preserving, it follows that  $\Box c \leq \Box(a \vee c)$  and  $\Box c \leq \Box(b \vee c)$ . Then, we have

$$\begin{aligned} \Box m(a, b, c) &= \Box[(a \vee c) \wedge (b \vee c)] = \Box(a \vee c) \wedge \Box(b \vee c) \\ &= (\Box(a \vee c) \vee \Box c) \wedge (\Box(b \vee c) \vee \Box c) = m(\Box(a \vee c), \Box(b \vee c), \Box c). \end{aligned}$$

Hence, identity (M) holds in  $A$ . Now, conversely, suppose that (M) holds in  $A$ . We need to check that condition (2) in Definition 4.4 is satisfied. First, we show that  $\Box$  is order-preserving. Let  $a, b \in A$  be such that  $a \leq b$ . So  $b = b \vee a = m(b, b, a)$ . Then, we have

$$\Box b = \Box m(b, b, a) = m(\Box(b \vee a), \Box(b \vee a), \Box a) = m(\Box b, \Box b, \Box a) = \Box b \vee \Box a.$$

Thus we obtain that  $\Box a \leq \Box b$ . Now let  $a, b \in A$  be such that  $a \wedge b$  there exists. Since  $\Box$  is order-preserving, we have  $\Box(a \wedge b) \leq \Box a, \Box b$ . Hence

$$\begin{aligned} \Box(a \wedge b) &= \Box m(a, b, a \wedge b) = m(\Box a, \Box b, \Box(a \wedge b)) \\ &= (\Box a \vee \Box(a \wedge b)) \wedge (\Box b \vee \Box(a \wedge b)) = \Box a \wedge \Box b. \end{aligned}$$

Therefore  $\langle A, m, \Box, 1 \rangle$  is a  $\Box$ -modal distributive nearlattice. □

Consider the DN-based logic  $\mathcal{S}_{\Box\mathbb{DN}}$  on the algebraic language  $\mathcal{L} = \{m, \Box, \top\}$  defined by (3.4) and (3.5). We already know that  $\mathcal{S}_{\Box\mathbb{DN}}$  satisfies properties (A1)-(A4). Moreover, it is straightforward to show directly that the following conditions hold:

- (N)  $\vdash_{\mathcal{S}_{\Box\mathbb{DN}}} \varphi$  implies  $\vdash_{\mathcal{S}_{\Box\mathbb{DN}}} \Box\varphi$ ;
- ( $\Box m$ )  $\Box m(\varphi, \psi, \chi) \vdash_{\mathcal{S}_{\Box\mathbb{DN}}} m(\Box(\varphi \vee \chi), \Box(\psi \vee \chi), \Box\chi)$ ;
- ( $m\Box$ )  $m(\Box(\varphi \vee \chi), \Box(\psi \vee \chi), \Box\chi) \vdash_{\mathcal{S}_{\Box\mathbb{DN}}} \Box m(\varphi, \psi, \chi)$ .

Moreover, notice that  $\mathbf{K}_{\mathcal{S}_{\Box\mathbb{DN}}} = \text{Alg}(\mathcal{S}_{\Box\mathbb{DN}}) = \Box\mathbb{DN}$  (see on page 12). Now let us show that  $\mathcal{S}_{\Box\mathbb{DN}}$  is the weakest selfextensional non-pseudo axiomatic logic satisfying conditions (A1)-(A4), (N), ( $\Box m$ ) and ( $m\Box$ ).

**Proposition 4.6** *Let  $\mathcal{S}$  be a selfextensional non-pseudo axiomatic logic satisfying conditions (B1)-(A4), (N), ( $\Box m$ ) and ( $m\Box$ ). Then,  $\mathcal{S}$  is an extension of  $\mathcal{S}_{\Box\mathbb{DN}}$ .*

*Proof* Let  $\mathcal{S}$  be a selfextensional non-pseudo axiomatic logic satisfying conditions (A1)-(A4), (N), ( $\Box m$ ) and ( $m\Box$ ). Then, by Theorems 3.8 and 3.10, we have that  $\mathcal{S}$  is DN-based relative to  $\text{Alg}(\mathcal{S}) = \mathbf{K}_{\mathcal{S}}$ . Since  $\mathcal{S}$  satisfies conditions (N), ( $\Box m$ ) and ( $m\Box$ ), it is straightforward to show that  $\mathbf{K}_{\mathcal{S}} \subseteq \Box\mathbb{DN}$ . Hence, since  $\mathbf{K}_{\mathcal{S}} \subseteq \Box\mathbb{DN} = \mathbf{K}_{\mathcal{S}_{\Box\mathbb{DN}}}$ , we obtain by Theorem 3.14 that  $\mathcal{S}_{\Box\mathbb{DN}} \leq \mathcal{S}$ . □

### 5 The logic preserving truth and the logic preserving degrees of truth

Let  $\mathcal{S}$  be a selfextensional logic with a DN-term  $m$ . We know that the canonical class of algebras  $\text{Alg}(\mathcal{S})$  associated with  $\mathcal{S}$  is a DN-based variety, that is,  $\text{Alg}(\mathcal{S})$  is a variety and for every  $A \in \text{Alg}(\mathcal{S})$ , the  $\{m\}$ -reduct  $\langle A, m^A \rangle$  is a distributive nearlattice. Hence,

every algebra  $A$  in  $\text{Alg}(\mathcal{S})$  has associated a partial order. Thus, a sentential logic can be defined using this class of algebras as follows: a formula  $\varphi$  is a logical consequence from some premises if and only if for every algebra and every interpretation, whenever the interpretation of the premises have a common lower bound, the interpretation of the formula  $\varphi$  also has the same lower bound; this logic is the so-called *the logic that preserves degrees of truth*.

Let now  $\mathcal{S}$  be a selfextensional logic with a DN-term  $m$  and with theorems. Thus, for every  $A \in \text{Alg}(\mathcal{S})$ , the distributive nearlattice  $\langle A, m^A \rangle$  has a greatest element  $h\varphi$ , where  $\varphi$  is any theorem of  $\mathcal{S}$  and  $h$  is any interpretation on  $A$ . We denote this element, for every algebra  $A$ , by  $1^A$ . If for every algebra  $A$  we consider that  $1^A$  is the only designated truth-value (*the truth*), then it can be defined a logic as follows: a formula  $\varphi$  follows logically from some premises if and only if for every algebra and every interpretation, whenever the interpretation of the premises are true, then the interpretation of the formula  $\varphi$  is true. This logic is known in the literature as *the logic preserving truth* (or as *the assertional logic*) associated with  $\text{Alg}(\mathcal{S})$ .

In this section, given a selfextensional logic  $\mathcal{S}$  with a DN-term (and with theorems), we study the connections between this logic  $\mathcal{S}$  and both the logic that preserves degrees of truth and the logic preserving truth associated with the class of algebras  $\text{Alg}(\mathcal{S})$ .

### 5.1 The logic preserving degrees of truth

*Logics preserving degrees of truth* associated with some particular ordered algebraic structures are studied and discussed in several articles, for instance see [20,21,24,37]. In particular, [20] is an interesting contribution to the discussion on the role of degrees of truth in many-valued logics from the point of view of Abstract Algebraic Logic.

Let  $\mathbf{K}$  be a class of algebras such that every algebra  $A \in \mathbf{K}$  has a partial order  $\leq$  associated with its universe. Thus, *the logic preserving degrees of truth* with respect to  $\mathbf{K}$  is defined as follows: let  $\varphi_1, \dots, \varphi_n, \varphi \in Fm$ ,

$$\varphi_1, \dots, \varphi_n \vdash_{\mathbf{K}}^{\leq} \varphi \iff (\forall A \in \mathbf{K})(\forall h \in \text{Hom}(Fm, A)) \quad (5.1)$$

$$(\forall a \in A)(\text{if } a \leq h\varphi_i \ \forall i = 1, \dots, n, \text{ then } a \leq h\varphi)$$

and

$$\emptyset \vdash_{\mathbf{K}}^{\leq} \varphi \iff (\forall A \in \mathbf{K})(\forall h \in \text{Hom}(Fm, A))(\forall a \in A)(a \leq h\varphi). \quad (5.2)$$

For an arbitrary  $\Gamma \subseteq Fm$  and  $\varphi \in Fm$  we define

$$\Gamma \vdash_{\mathbf{K}}^{\leq} \varphi \iff \text{there is a finite } \Gamma_0 \subseteq \Gamma \text{ such that } \Gamma_0 \vdash_{\mathbf{K}}^{\leq} \varphi. \quad (5.3)$$

**Definition 5.1** Let  $\mathbf{K}$  be a DN-based class of algebras. The *logic preserving degrees of truth with respect to  $\mathbf{K}$* ,  $\mathcal{S}_{\mathbf{K}}^{\leq} = \langle Fm, \vdash_{\mathbf{K}}^{\leq} \rangle$ , is defined by (5.1)–(5.3).

**Proposition 5.2** *The logic  $\mathcal{S}_{\mathbf{K}}^{\leq}$  is an extension of the logic  $\mathcal{S}_{\mathbf{K}}$ . Moreover,  $\mathcal{S}_{\mathbf{K}}$  and  $\mathcal{S}_{\mathbf{K}}^{\leq}$  have the same theorems, that is,  $\vdash_{\mathbf{K}} \varphi$  if and only if  $\vdash_{\mathbf{K}}^{\leq} \varphi$ .*




Fig. 2 The distributive nearlattice of Example 5.4

Let  $K$  be a DN-based class of algebras. From (5.1), it is clear that for all  $\varphi, \psi \in Fm$ ,

$$\varphi \vdash_K^{\leq} \psi \iff (\forall A \in K)(\forall h \in \text{Hom}(Fm, A))(h\varphi \leq h\psi).$$

Thus, the following proposition is straightforward. Recall that for a logic  $S$ ,  $K_S$  denotes the intrinsic variety of  $S$ .

**Proposition 5.3** *Let  $K$  be a DN-based class. Then,*

1. *the logic  $S_K^{\leq}$  is selfextensional;*
2.  *$\mathbb{V}(K) = K_{S_K^{\leq}}$ ;*
3.  *$S_K^{\leq}$  satisfies properties (A1)-(A3).*

*Example 5.4* The logic  $S_K^{\leq}$  does not necessarily satisfy property (A4), and thus the logics  $S_K^{\leq}$  and  $S_K$  may be different. Consider the language  $\mathcal{L} := \{m, \perp_1, \perp_2, \top\}$  of type  $(3, 0, 0, 0)$ . Let  $A$  be the distributive nearlattice given in Fig. 2, and such that  $\perp_1^A = a$ ,  $\perp_2^A = b$  and  $\top^A = 1$ . Let  $x$  be an arbitrary variable. Then, we have  $\perp_1, \perp_2 \vdash_A^{\leq} x$  but  $m(\perp_1, \perp_2, x) \not\vdash_A^{\leq} x$ . Hence,  $S_{\{A\}}^{\leq}$  does not satisfy property (A4). Moreover, notice that  $S_{\{A\}}^{\leq}$  is not a DN-based logic. Otherwise, it would satisfy (A4).

**Proposition 5.5** *Let  $K$  be a DN-based class. Then, the logic  $S_K^{\leq}$  satisfies property (A4) if and only if  $S_K = S_K^{\leq}$ .*

*Proof* Assume that  $S_K^{\leq}$  satisfies property (A4). By Proposition 5.2 we need only to prove that  $S_K^{\leq} \leq S_K$ . Let  $\varphi_0, \dots, \varphi_n, \varphi$  be formulas and assume that  $\varphi_0, \dots, \varphi_n \vdash_K^{\leq} \varphi$ . Thus, by (A4),  $m^n(\varphi_0, \dots, \varphi_n, \varphi) \vdash_K^{\leq} \varphi$ . Then, it is straightforward to check that  $m^n(\varphi_0, \dots, \varphi_n, \varphi) \vdash_K \varphi$ . Now, by condition (3) of Proposition 3.2, it follows that  $\varphi_0, \dots, \varphi_n \vdash_K \varphi$ . Hence  $S_K^{\leq} \leq S_K$ . Conversely, if  $S_K = S_K^{\leq}$ , then it is clear that  $S_K^{\leq}$  satisfies property (A4).  $\square$

**Proposition 5.6** *If  $K$  is a DN-based variety, then*

$$K_{S_K^{\leq}} = K = K_{S_K} = \text{Alg}(S_K).$$

*Proof* It follows from Propositions 5.3 and 3.12, and Theorem 3.10.  $\square$

For the next results, we need the following concepts. Let  $P$  be a partially ordered set. Let  $X \subseteq P$ . We set  $X^\ell := \{a \in P : a \leq x, \text{ for all } x \in X\}$  and  $X^u := \{a \in P : a \geq x, \text{ for all } x \in X\}$ . A subset  $F \subseteq P$  is said to be a *Frink filter* [28] if for every finite  $X \subseteq F$ , we have  $X^{\ell u} \subseteq F$ . We denote by  $\text{Fi}_F(P)$  the collection of all Frink filters of  $P$ . It is easy to check that  $\text{Fi}_F(P)$  is an algebraic closure system on  $P$ .

Then, condition (5.1) can be written as

$$\varphi_1, \dots, \varphi_n \vdash_{\bar{K}} \varphi \iff (\forall A \in \mathbf{K})(\forall h \in \text{Hom}(Fm, A)) \\ h\varphi \in \{h\varphi_1, \dots, h\varphi_n\}^{\ell u}.$$

The next lemma is straightforward, and thus we omit its proof.

**Lemma 5.7** *For every DN-class of algebras  $\mathbf{K}$  and every algebra  $A \in \mathbf{K}$ , we have  $\text{Fi}_F(A) \subseteq \text{Fi}_{\mathcal{S}_{\bar{K}}^\leq}(A)$ .*

**Proposition 5.8** *Let  $\mathbf{K}$  be a DN-based variety. Then, we have*

$$\text{Alg}(\mathcal{S}_{\bar{K}}^\leq) = \mathbf{K} = \text{Alg}(\mathcal{S}_{\mathbf{K}}).$$

*Proof* By Lemma 2.3, we know that  $\text{Alg}(\mathcal{S}_{\bar{K}}^\leq) \subseteq \mathbf{K}_{\mathcal{S}_{\bar{K}}^\leq} = \mathbf{K}$ . Recall that  $A \in \text{Alg}(\mathcal{S}_{\bar{K}}^\leq)$  if and only if there is an algebraic closure system  $\mathcal{C}$  on  $A$  such that  $\langle A, \mathcal{C} \rangle \in \mathbf{GMod}^*(\mathcal{S}_{\bar{K}}^\leq)$ . Let  $A \in \mathbf{K}$ . Since  $\text{Fi}_F(A)$  is an algebraic closure system and  $\text{Fi}_F(A) \subseteq \text{Fi}_{\mathcal{S}_{\bar{K}}^\leq}(A)$ , it follows that  $\langle A, \text{Fi}_F(A) \rangle \in \mathbf{GMod}^*(\mathcal{S}_{\bar{K}}^\leq)$ .

Notice that the upsets  $[a] = \{x \in A : a \leq x\}$  are Frink filters of  $A$ . So, we have  $\langle a, b \rangle \in \Lambda_A(\text{Fi}_F(A))$  if and only if  $a = b$ . Then, the Frege relation  $\Lambda_A(\text{Fi}_F(A))$  of the g-model  $\langle A, \text{Fi}_F(A) \rangle$  is the identity relation. Hence,  $\langle A, \text{Fi}_F(A) \rangle \in \mathbf{GMod}^*(\mathcal{S}_{\bar{K}}^\leq)$ ; and thus,  $A \in \text{Alg}(\mathcal{S}_{\bar{K}}^\leq)$ . Therefore, by Proposition 5.6,  $\text{Alg}(\mathcal{S}_{\bar{K}}^\leq) = \mathbf{K} = \text{Alg}(\mathcal{S}_{\mathbf{K}})$ .  $\square$

The following corollary is a consequence of Lemma 5.7.

**Corollary 5.9** *Let  $\mathbf{K}$  be a DN-based variety. Then, the logic  $\mathcal{S}_{\bar{K}}^\leq$  is fully selfextensional.*

Let  $\mathbf{K}$  be a DN-based class of algebras and  $A \in \mathbf{K}$ . Since  $\text{Fi}_F(A)$  is a closure system,  $\text{Fi}_F(A)$  is a lattice. Let us show that the distributivity of the lattices  $\text{Fi}_F(A)$  is a sufficient condition for the logics  $\mathcal{S}_{\mathbf{K}}$  and  $\mathcal{S}_{\bar{K}}^\leq$  to coincide.

**Proposition 5.10** *Let  $\mathbf{K}$  be a DN-based variety. If for every  $A \in \mathbf{K}$  the lattice  $\text{Fi}_F(A)$  is distributive, then  $\mathcal{S}_{\mathbf{K}} = \mathcal{S}_{\bar{K}}^\leq$ .*

*Proof* Assume that for each  $A \in \mathbf{K}$ , the lattice  $\text{Fi}_F(A)$  is distributive. Since the logic  $\mathcal{S}_{\bar{K}}^\leq$  is an extension of the logic  $\mathcal{S}_{\mathbf{K}}$ , we have  $\text{Fi}_{\mathcal{S}_{\bar{K}}^\leq}(A) \subseteq \text{Fi}_{\mathcal{S}_{\mathbf{K}}}(A)$  for every algebra  $A$ . Let now  $A \in \mathbf{K}$ . Since the lattice  $\text{Fi}_F(A)$  is distributive, it follows that  $\text{Fi}_F(A) = \text{Fi}(A)$  (see [30, Proposition 4.3]). Then, by Proposition 3.9 and Lemma 5.7, it follows that

$$\text{Fi}_{\mathcal{S}_{\mathbf{K}}}(A) \setminus \{\emptyset\} = \text{Fi}(A) = \text{Fi}_F(A) \subseteq \text{Fi}_{\mathcal{S}_{\bar{K}}^\leq}(A). \tag{5.4}$$

Hence, since  $\mathcal{S}_K$  and  $\mathcal{S}_K^{\leq}$  have the same theorems, we obtain  $\text{Fi}_{\mathcal{S}_K}(A) = \text{Fi}_{\mathcal{S}_K^{\leq}}(A)$  for all  $A \in K$ . Now, notice that every sentential logic  $\mathcal{S}$  is complete with respect to the class of g-models  $\{(A, \text{Fi}_{\mathcal{S}}(A)) : A \in \text{Alg}(\mathcal{S})\}$ . Thus, by Proposition 5.8, we obtain that  $\mathcal{S}_K = \mathcal{S}_K^{\leq}$ .  $\square$

*Remark 5.11* Let  $K$  be a DN-based variety such that the lattice  $\text{Fi}_{\mathcal{F}}(A)$  is distributive for every  $A \in K$ . Then, the proof of Proposition 5.10 (see (5.4)) shows that if the logic  $\mathcal{S}_K$  has theorems, then  $\text{Fi}_{\mathcal{S}_K}(A) = \text{Fi}_{\mathcal{F}}(A)$  for all  $A \in K$ .

### 5.2 The logic preserving truth

We will say that a class of algebras  $K$  is *DN<sup>1</sup>-based* if it is DN-based and every algebra  $A$  in  $K$  has a greatest element, that is, there is  $1^A \in A$  such that  $a \leq 1^A$  for all  $a \in A$ . Notice that if  $\mathcal{S}$  is a selfextensional logic with a DN-term and with theorems, then  $\text{Alg}(\mathcal{S})$  is a DN<sup>1</sup>-based variety and for every  $A \in \text{Alg}(\mathcal{S})$ ,  $1^A = h\varphi$  for any theorem  $\varphi$  and any interpretation  $h$  on  $A$ . Moreover, since  $\mathcal{S}$  has a theorem, it follows by Proposition 3.12 that  $\mathcal{S} = \mathcal{S}_{\text{Alg}(\mathcal{S})}$ . From now on, unless otherwise stated,  $K$  will denote a DN<sup>1</sup>-based variety.

**Definition 5.12** The *logic preserving truth associated with  $K$* , denoted by  $\mathcal{S}_K^1 = \langle Fm, \vdash_K^1 \rangle$ , is defined as follows: let  $\Gamma \cup \{\varphi\} \subseteq Fm$  be finite,

$$\Gamma \vdash_K^1 \varphi \iff (\forall A \in K)(\forall h \in \text{Hom}(Fm, A)) \quad (5.5)$$

$$(h\psi = 1^A \text{ for all } \psi \in \Gamma \implies h\varphi = 1^A).$$

For an arbitrary  $\Gamma \subseteq Fm$ ,  $\Gamma \vdash_K^1 \varphi$  if and only if there is a finite  $\Gamma_0 \subseteq \Gamma$  such that  $\Gamma_0 \vdash_K^1 \varphi$ .

**Proposition 5.13** The logic  $\mathcal{S}_K^1$  is an extension of  $\mathcal{S}_K^{\leq}$ , and hence it is an extension of the logic  $\mathcal{S}_K$ . Moreover, the logics  $\mathcal{S}_K$ ,  $\mathcal{S}_K^{\leq}$  and  $\mathcal{S}_K^1$  have the same theorems.

*Example 5.14* Let  $K$  be a variety of residuated lattices [29]. Then, it is clear that  $K$  is a DN-based variety. In [6] the authors prove (Theorem 4.12) that  $\vdash_K^1 = \vdash_K^{\leq}$  if and only if  $K$  is a variety of generalised Heyting algebras [38].<sup>2</sup> Then, by Example 5.4, we have for a DN-based class of algebras  $K$  that the logics  $\mathcal{S}_K$ ,  $\mathcal{S}_K^{\leq}$  and  $\mathcal{S}_K^1$  may be different.

Let  $A$  be an algebra and  $F \subseteq A$ . The *Leibniz congruence of  $F$  relative to  $A$* , denoted by  $\Omega^A F$ , is the greatest congruence of  $A$  compatible with  $F$ , that is, it does not relate elements in  $F$  with elements not in  $F$ . The mapping  $F \mapsto \Omega^A F$  is called the *Leibniz operator* of the algebra  $A$  and it is denoted by  $\Omega^A$ . This operator is an essential tool in Abstract Algebraic Logic for classifying sentential logics. The structure of this classification of sentential logics is called the *Leibniz hierarchy*. We address the reader to [17, 23, 27] for more information on this hierarchy.

<sup>2</sup> Generalised Heyting algebras are called relatively pseudo-complemented lattices in [38].

**Proposition 5.15** *If for every  $A \in \mathbf{K}$  and every nonempty and finite  $X \subseteq A$  we have  $\text{Fi}_A(X) = \{a \in A : \langle a, 1^A \rangle \in \Omega^A \text{Fi}_A(X)\}$ , then  $\mathcal{S}_{\mathbf{K}} = \mathcal{S}_{\mathbf{K}}^1$ .*

*Proof* By Proposition 5.13, we only need to prove that for all  $\varphi_0, \dots, \varphi_n, \varphi \in Fm$ ,  $\varphi_0, \dots, \varphi_n \vdash_{\mathbf{K}}^1 \varphi$  implies  $\varphi_0, \dots, \varphi_n \vdash_{\mathbf{K}} \varphi$ . So, assume that  $\varphi_0, \dots, \varphi_n \vdash_{\mathbf{K}}^1 \varphi$ . Let  $A \in \mathbf{K}$  and  $h \in \text{Hom}(Fm, A)$ . We set  $F := \text{Fi}_A(h\varphi_0, \dots, h\varphi_n)$ . Since  $\mathbf{K}$  is a variety and  $\Omega^A F$  is a congruence on  $A$ , it follows that  $A/\Omega^A F \in \mathbf{K}$ . By hypothesis, we have that  $F = 1^A/\Omega^A F = 1^{A/\Omega^A F}$ . Let now  $\widehat{h} := \pi \circ h: Fm \rightarrow A/\Omega^A F$ , where  $\pi: A \rightarrow A/\Omega^A F$  is the natural map. Thus,  $\widehat{h}\varphi_i = h\varphi_i/\Omega^A F = 1^A/\Omega^A F = 1^{A/\Omega^A F}$  for all  $i = 0, 1, \dots, n$ . Then,  $\widehat{h}\varphi = 1^{A/\Omega^A F}$ . Hence  $h\varphi \in F = \text{Fi}_A(h\varphi_0, \dots, h\varphi_n)$ . Thus, by Proposition 2.15, we obtain that  $h\varphi = m^n(h\varphi_0, \dots, h\varphi_n, h\varphi)$ . Hence,  $\varphi_0, \dots, \varphi_n \vdash_{\mathbf{K}} \varphi$ .  $\square$

One of the most important and large classes of sentential logics under the point of view of Abstract Algebraic Logic is the class of *protoalgebraic logics*. This class of logics was introduced and studied by Blok and Pigozzi [4], and independently it was considered by Czelakowski [16]. There are several useful characterizations of the notion of protoalgebraicity. For our purposes, we choose the following as the definition of protoalgebraic logic.

**Definition 5.16** A sentential logic  $\mathcal{S}$  is said to be *protoalgebraic* if there is a set of formulas in two variables  $\Delta(x, y)$  such that  $\emptyset \vdash_{\mathcal{S}} \Delta(x, x)$  and  $x, \Delta(x, y) \vdash_{\mathcal{S}} y$ . A set with these two properties will be called a *set of protoimplication formulas* for  $\mathcal{S}$ .

Since the logics in this paper are considered to be finitary, the sets of protoimplication formulas can always be chosen to be finite.

Algebraizable logics, finitely algebraizable logics and regularly algebraizable logics are important classes of protoalgebraic logics, see for instance [5, 17, 23, 27]. A sentential logic  $\mathcal{S}$  is *algebraizable* if and only if (i) there is a set of formulas in two variables  $\Delta(x, y)$  such that for each algebra  $A$  and each  $\mathcal{S}$ -filter  $F$  of  $A$ ,  $\Omega^A F = \{\langle a, b \rangle \in A^2 : \Delta^A(a, b) \subseteq F\}$ , and (ii) there is a set of equations  $\tau(x)$  in one variable such that for every algebra  $A$  and every  $\mathcal{S}$ -filter  $F$  of  $A$  with  $\Omega^A F$  being the identity relation,  $F = \{a \in A : A \models \tau(x)[a]\}$ . The set  $\Delta(x, y)$  is called a set of *equivalence formulas* for  $\mathcal{S}$ . It follows that every set of equivalence formulas is a set of protoimplication formulas.

A logic  $\mathcal{S}$  is said to be *finitely algebraizable* when it is algebraizable and the sets  $\Delta(x, y)$  and  $\tau(x)$  are finite. A logic  $\mathcal{S}$  is said to be *regularly algebraizable* if it is finitely algebraizable and for every set of equivalence formulas  $\Delta(x, y)$  the G-rule  $x, y \vdash_{\mathcal{S}} \Delta(x, y)$  is satisfied.

A quasivariety  $\mathbf{K}$  is said to be *pointed* if there is a term  $\varphi(x_1, \dots, x_n)$  with the property that  $\varphi(x_1, \dots, x_n) \approx \varphi(y_1, \dots, y_n)$  is valid in  $\mathbf{K}$  for all variables  $y_1, \dots, y_n$ . Such a term is called a *constant term* since it behaves like a constant. Once we fix a constant term, we will denote it by 1; we will say that  $\mathbf{K}$  is *1-pointed* and we will use  $1^A$  to refer the interpretation of the constant term 1 in  $A$ , for each  $A \in \mathbf{K}$ . Notice that if  $\mathcal{S}$  is a selfextensional logic with a DN-term and with theorems, then the variety  $\text{Alg}(\mathcal{S})$  is 1-pointed, where the constant term is any theorem of  $\mathcal{S}$ , and for each  $A \in \text{Alg}(\mathcal{S})$ ,  $1^A$  is the greatest element of  $\langle A, m^A \rangle$ .

A 1-pointed quasivariety  $K$  is said to be *relatively point regular* when for every  $A \in K$  and every  $\theta, \theta' \in \text{Con}_K(A)$ , if  $1^A/\theta = 1^A/\theta'$ , then  $\theta = \theta'$ . If  $K$  is a variety, then we say simply that  $K$  is *point regular*, since  $\text{Con}_K(A) = \text{Con}(A)$  for every  $A \in K$ .

**Theorem 5.17** [19] *A logic  $\mathcal{S}$  is regularly algebraizable if and only if  $\text{Alg}(\mathcal{S})$  is 1-pointed and relatively point regular quasivariety and  $\mathcal{S} = \mathcal{S}_{\text{Alg}(\mathcal{S})}^1$ .*

Let  $\langle A, m \rangle$  be a distributive nearlattice. Then,  $\theta$  is a congruence on  $A$  if and only if (i) if  $\langle a, b \rangle, \langle c, d \rangle \in \theta$ , then  $\langle a \vee c, b \vee d \rangle \in \theta$ , and (ii) if  $\langle a, b \rangle, \langle c, d \rangle \in \theta$  and  $a \wedge c, b \wedge d$  exist in  $A$ , then  $\langle a \wedge c, b \wedge d \rangle \in \theta$ , see [32].

**Theorem 5.18** *Let  $\mathcal{S}$  be a selfextensional logic with a DN-term and with theorems. If the logic  $\mathcal{S}$  is protoalgebraic, then  $\mathcal{S} = \mathcal{S}_{\text{Alg}(\mathcal{S})}^1$  if and only if  $\mathcal{S}$  is regularly algebraizable.*

*Proof* Let  $\mathcal{S}$  be a selfextensional logic with a DN-term and with theorems, and assume that  $\mathcal{S}$  is protoalgebraic. Recall that  $\mathcal{S} = \mathcal{S}_{\text{Alg}(\mathcal{S})}$ , and by Proposition 5.13 we have that  $\mathcal{S} \subseteq \mathcal{S}_{\text{Alg}(\mathcal{S})}^1$ . If  $\mathcal{S}$  is regularly algebraizable, then by the previous theorem we have that  $\mathcal{S} = \mathcal{S}_{\text{Alg}(\mathcal{S})}^1$ .

Now, conversely, assume that  $\mathcal{S} = \mathcal{S}_{\text{Alg}(\mathcal{S})}^1$ . We already know that  $\text{Alg}(\mathcal{S})$  is 1-pointed. To show that the logic  $\mathcal{S}$  is regularly algebraizable we only need to prove, by the previous theorem, that the variety  $\text{Alg}(\mathcal{S})$  is point regular. Let  $A \in \text{Alg}(\mathcal{S})$  and let  $\theta, \theta' \in \text{Con}(A)$  be such that  $1^A/\theta = 1^A/\theta'$ . As  $\mathcal{S}$  is finitary and protoalgebraic, let  $\Delta(x, y)$  be a finite set of protoimplication formulas for  $\mathcal{S}$ . Moreover, since the logic  $\mathcal{S}$  has theorems,  $\Delta(x, y)$  can be assumed to be nonempty. We set  $\Delta(x, y) = \{\varphi_1(x, y), \dots, \varphi_n(x, y)\}$ . Let  $\langle a, b \rangle \in \theta$ . So,

$$\{(\varphi^A(a, a), \varphi^A(a, b)) : \varphi(x, y) \in \Delta(x, y)\} \subseteq \theta. \tag{5.6}$$

Since  $\vdash_{\mathcal{S}} \Delta(x, x)$ , it follows that  $\{(1^A, \varphi^A(a, b)) : \varphi(x, y) \in \Delta(x, y)\} \subseteq \theta$ . Thus,  $\{(1^A, \varphi^A(a, b) \vee b) : \varphi(x, y) \in \Delta(x, y)\} \subseteq \theta$ . Then, by hypothesis,  $\{(1^A, \varphi^A(a, b) \vee b) : \varphi(x, y) \in \Delta(x, y)\} \subseteq \theta'$ . It follows that

$$(1^A, (\varphi_1^A(a, b) \vee b) \wedge \dots \wedge (\varphi_n^A(a, b) \vee b)) \in \theta'.$$

Then, we obtain

$$\langle a \vee b, (\varphi_1^A(a, b) \vee b) \wedge \dots \wedge (\varphi_n^A(a, b) \vee b) \wedge (a \vee b) \rangle \in \theta'.$$

Notice that

$$(\varphi_1^A(a, b) \vee b) \wedge \dots \wedge (\varphi_n^A(a, b) \vee b) \wedge (a \vee b) = m^n(\varphi_1^A(a, b), \dots, \varphi_n^A(a, b), a, b).$$

Thus,

$$\langle a \vee b, m^n(\varphi_1^A(a, b), \dots, \varphi_n^A(a, b), a, b) \rangle \in \theta'. \tag{5.7}$$



Now, since  $x, \Delta(x, y) \vdash_{\mathcal{S}} y$ , it follows by property (A4) that

$$m^n(\varphi_1(x, y), \dots, \varphi_n(x, y), x, y) \vdash_{\mathcal{S}} y.$$

Let  $h \in \text{Hom}(Fm, A)$  be such that  $h(x) = a$  and  $h(y) = b$ . Then, we obtain that

$$m^n(\varphi_1^A(a, b), \dots, \varphi_n^A(a, b), a, b) = b.$$

Thus, by (5.7), we have  $\langle a \vee b, b \rangle \in \theta'$ . With a similar argumentation we can get  $\langle a \vee b, a \rangle \in \theta'$ . Hence,  $\langle a, b \rangle \in \theta'$ . We have proved that  $\theta \subseteq \theta'$ . Similarly, we have  $\theta' \subseteq \theta$ . Then  $\theta = \theta'$ . Hence,  $\text{Alg}(\mathcal{S})$  is point regular. Therefore, since  $\text{Alg}(\mathcal{S})$  is 1-pointed and point regular and since  $\mathcal{S} = \mathcal{S}_{\text{Alg}(\mathcal{S})}^1$ , it follows by the previous theorem that  $\mathcal{S}$  is regularly algebraizable.  $\square$

### Conclusions

Given an algebraic language  $\mathcal{L}$  with a ternary term  $m$ , we have defined when  $m$  is a distributive nearlattice term (DN-term) for a sentential logic  $\mathcal{S}$  (see Definition 3.1). The term  $m$  is a DN-term for  $\mathcal{S}$  if it satisfies some syntactical properties ((A1)–(A4)); roughly speaking, these properties (A1)–(A4) mean that when  $m$  is interpreted in every algebra  $A$  of the algebraic counterpart of  $\mathcal{S}$  the  $\{m\}$ -reduct  $\langle A, m^A \rangle$  is a distributive nearlattice.

Then, we characterised the selfextensional logics with a DN-term  $m$  as those for which the consequence relation can be defined by the order induced by the ternary term  $m$  interpreted in the algebras of the algebraic counterpart of the logic (see Theorems 3.8 and 3.10).

Given a DN-based variety  $\mathbf{K}$ , we define the logic  $\mathcal{S}_{\mathbf{K}}$  (see on page 12); this logic is selfextensional with a DN-term, and  $\mathbf{K} = \text{Alg}(\mathcal{S}_{\mathbf{K}})$ . Since the algebras of  $\mathbf{K}$  have associated a partial order, it can be defined the logic preserving degrees of truth  $\mathcal{S}_{\mathbf{K}}^{\leq}$  (see Definition 5.1). We have shown some properties of the logic  $\mathcal{S}_{\mathbf{K}}^{\leq}$ . We found some sufficient conditions for logics  $\mathcal{S}_{\mathbf{K}}$  and  $\mathcal{S}_{\mathbf{K}}^{\leq}$  coincide (see Propositions 5.5 and 5.10).

If  $\mathcal{S}$  is a selfextensional logic with a DN-term and with theorems, then the algebraic counterpart  $\text{Alg}(\mathcal{S})$  of  $\mathcal{S}$  is 1-pointed. Thus, we can define the logic preserving truth  $\mathcal{S}_{\text{Alg}(\mathcal{S})}^1$  associated with  $\text{Alg}(\mathcal{S})$ . We also have found some sufficient conditions for logics  $\mathcal{S}$  and  $\mathcal{S}_{\text{Alg}(\mathcal{S})}^1$  coincide (see Proposition 5.15 and Theorem 5.18).

Selfextensional logics having a DN-term (or equivalently, DN-based logics) include those sentential logics defined by varieties of distributive lattices. More precisely, consider an algebraic language  $\mathcal{L}$  with two binary terms  $\wedge$  and  $\vee$  and let  $\mathbf{K}$  be a variety of type  $\mathcal{L}$  such that for each  $A \in \mathbf{K}$ ,  $\langle A, \wedge^A, \vee^A \rangle$  is a distributive lattice. Then, it is clear that  $\mathbf{K}$  is, in particular, a DN-based variety. On the one hand, since the algebras of  $\mathbf{K}$  have a conjunction, it can be defined in a natural way a logic  $\mathcal{S}'_{\mathbf{K}}$  as in [34, pp. 79] which is the semilattice-based logic associated with  $\mathbf{K}$ . On the other hand, since  $\mathbf{K}$  is a DN-based variety, we can consider the DN-based logic  $\mathcal{S}_{\mathbf{K}}$  associated with  $\mathbf{K}$  (see (3.4) and (3.5)). Then, it is straightforward to check that the two logics  $\mathcal{S}'_{\mathbf{K}}$  and  $\mathcal{S}_{\mathbf{K}}$  coincide.

Let  $K$  be a  $DN^1$ -based variety (see Sect. 5.2). Since distributive nearlattices can be considered as generalisations both of Tarski algebras and of distributive lattices, it will be important to carry out studies on the logics  $S_K$ ,  $S_K^{\leq}$  and  $S_K^1$  under the point of view of AAL. In particular, we will study how these logics behave concerning the classifications in the hierarchy of Leibniz and the hierarchy of Frege. These task will be pursued elsewhere. The papers [6, 22, 25, 35] will be of great help to carry out these investigations.

**Acknowledgements** I would like to thank Prof. Ramon Jansana for his valuable comments and suggestions on an earlier version of the manuscript that helped to improve the presentation. I would also like to thank the anonymous referee, whose comments and suggestions helped to improve the paper.

## References

1. Abbott, J.C.: Implicational algebras. *Bull. Math. R. S. Roumaine* **11**(1), 3–23 (1967)
2. Abbott, J.C.: Semi-boolean algebra. *Matematički Vesnik* **4**(40), 177–198 (1967)
3. Araújo, J., Kinyon, M.: Independent axiom systems for nearlattices. *Czech. Math. J.* **61**(4), 975–992 (2011)
4. Blok, W.J., Pigozzi, D.: Protoalgebraic logics. *Stud. Logica* **45**(4), 337–369 (1986)
5. Blok, W.J., Pigozzi, D.: *Algebraizable Logics*, vol. 77. American Mathematical Society, Providence (1989)
6. Bou, F., Esteva, F., Font, J.M., Gil, A., Godo, L., Torrens, A., Verdú, V.: Logics preserving degrees of truth from varieties of residuated lattices. *J. Logic Comput.* **19**(6), 1031–1069 (2009)
7. Calomino, I.: Supremo álgebra distributivas: una generalización de las álgebra de Tarski. Ph.D. thesis, Universidad Nacional del Sur (2015)
8. Celani, S., Calomino, I.: Stone style duality for distributive nearlattices. *Algebra Univ.* **71**(2), 127–153 (2014)
9. Celani, S., Calomino, I.: On homomorphic images and the free distributive lattice extension of a distributive nearlattice. *Rep. Math. Logic* **51**, 57–73 (2016)
10. Chajda, I., Halaš, R., Kühr, J.: *Semilattice Structures*, vol. 30. Heldermann, Lemgo (2007)
11. Chajda, I., Halaš, R.: An example of a congruence distributive variety having no near-unanimity term. *Acta Univ. M. Belii Ser. Math.* **13**, 29–31 (2006)
12. Chajda, I., Kolařík, M.: A decomposition of homomorphic images of nearlattices. *Acta Univ. Palacki. Olomuc. Fac. rer. nat. Mathematica* **45**(1), 43–51 (2006)
13. Chajda, I., Kolařík, M.: Nearlattices. *Discrete Math.* **308**(21), 4906–4913 (2008)
14. Cintula, P., Noguera, C.: The proof by cases property and its variants in structural consequence relations. *Studia Logica* 1–35 (2011)
15. Cornish, W., Hickman, R.C.: Weakly distributive semilattices. *Acta Math. Hungar.* **32**(1), 5–16 (1978)
16. Czelakowski, J.: Algebraic aspects of deduction theorems. *Stud. Log.* **44**(4), 369–387 (1985)
17. Czelakowski, J.: *Protoalgebraic Logics*. Kluwer Academic Publishers, Dordrecht (2001)
18. Czelakowski, J.: The Suszko operator. Part I. *Stud. Log.* **74**(1–2), 181–231 (2003)
19. Czelakowski, J., Pigozzi, D.: Fregean logics. *Ann. Pure Appl. Log.* **127**(1–3), 17–76 (2004)
20. Font, J.M.: An Abstract Algebraic Logic View of Some Multiple-valued Logics, pp. 25–57. *Physica-Verlag HD, Heidelberg* (2003). [https://doi.org/10.1007/978-3-7908-1769-0\\_2](https://doi.org/10.1007/978-3-7908-1769-0_2)
21. Font, J.M.: Taking degrees of truth seriously. *Stud. Log.* **91**(3), 383–406 (2009)
22. Font, J.M.: On semilattice-based logics with an algebraizable assertional companion. *Rep. Math. Log.* **46**, 109–132 (2011)
23. Font, J.M.: *Abstract Algebraic Logic—An Introductory Textbook*, *Studies in Logic*, vol. 60. College Publications, London (2016)
24. Font, J.M., Gil, A., Torrens, A., Verdú, V.: On the infinite-valued Łukasiewicz logic that preserves degrees of truth. *Arch. Math. Log.* **45**(7), 839–868 (2006)
25. Font, J.M., Jansana, R.: Leibniz filters and the strong version of a protoalgebraic logic. *Arch. Math. Logic* **40**(6), 437–465 (2001)

26. Font, J.M., Jansana, R.: A General Algebraic Semantics for Sentential Logics. Lecture Notes in Logic, vol. 7, 2nd edn. The Association for Symbolic Logic, Poughkeepsie (2009)
27. Font, J.M., Jansana, R., Pigozzi, D.: A survey of abstract algebraic logic. *Stud. Log.* **74**(1–2), 13–97 (2003)
28. Frink, O.: Ideals in partially ordered sets. *Am. Math. Mon.* **61**(4), 223–234 (1954)
29. Galatos, N., Jipsen, P., Kowalski, T., Ono, H.: *Residuated Lattices: An Algebraic Glimpse at Substructural Logics*. Elsevier, New York (2007)
30. González, L.: The Free Distributive Semilattice Extension of a Distributive Poset (2017)
31. González, L.: The logic of distributive nearlattices. *Soft. Comput.* **9**(22), 2797–2807 (2018). <https://doi.org/10.1007/s00500-017-2750-0>
32. Hickman, R.: Join algebras. *Commun Algebra* **8**(17), 1653–1685 (1980)
33. Jansana, R.: Selfextensional logics with implication. In: Beziau, J.Y. (ed.) *Logica Universalis*, pp. 65–88. Springer, Berlin (2005)
34. Jansana, R.: Selfextensional logics with a conjunction. *Stud. Log.* **84**(1), 63–104 (2006)
35. Jansana, R.: Algebraizable logics with a strong conjunction and their semi-lattice based companions. *Arch. Math. Log.* **51**(7), 831–861 (2012). <https://doi.org/10.1007/s00153-012-0301-z>
36. Łoś, J., Suszko, R.: Remarks on sentential logics. *Indag. Math.* **20**, 177–183 (1958)
37. Nowak, M.: Logics preserving degrees of truth. *Stud. Log.* **49**(4), 483–499 (1990)
38. Rasiowa, H.: *An Algebraic Approach to Non-Classical Logics*. North-Holland (1974)
39. Rebagliato, J., Verdú, V.: Algebraizable Gentzen Systems and the Deduction Theorem for Gentzen Systems, vol. 175, Mathematics Preprint Series (1995)
40. Wang, S., Cintula, P.: Logics with disjunction and proof by cases. *Arch. Math. Log.* **47**(5), 435–446 (2008)